

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

Genome mapping of quantitative trait loci (QTL) controlling domestication traits of intermediate wheatgrass (*Thinopyrum intermedium*).

### Permalink

<https://escholarship.org/uc/item/67b4r6bd>

### Journal

TAG. Theoretical and applied genetics. Theoretische und angewandte Genetik, 132(8)

### ISSN

0040-5752

### Authors

Larson, Steve  
DeHaan, Lee  
Poland, Jesse  
et al.

### Publication Date

2019-08-01

### DOI

10.1007/s00122-019-03357-6

Peer reviewed

**Title page**

**Genome mapping of quantitative trait loci (QTL) controlling domestication traits of intermediate wheatgrass (*Thinopyrum intermedium*)**

Steve Larson<sup>1\*</sup>, Lee DeHaan<sup>2</sup>, Jesse Poland<sup>3</sup>, Xiaofei Zhang<sup>4</sup>, Kevin Dorn<sup>3</sup>, Traci Kantarski<sup>5</sup>, James Anderson<sup>6</sup>, Jeremy Schmutz<sup>7,8</sup>, Jane Grimwood<sup>8</sup>, Jerry Jenkins<sup>8</sup>, Shengqiang Shu<sup>7</sup>, Jared Crain<sup>3</sup>, Matthew Robbins<sup>1</sup>, Kevin Jensen<sup>1</sup>

<sup>1</sup>United States Department of Agriculture, Agriculture Research Service, Forage and Range Research, Utah State University, Logan, UT 84322, United States

<sup>2</sup>The Land Institute, 2440 E. Water Well Rd, Salina, KS 67401, United States

<sup>3</sup>Department of Plant Pathology, Kansas State University, 4024 Throckmorton, Manhattan, KS 66506, United States

<sup>4</sup>Department of Horticultural Science, North Carolina State University, 212 Kilgore Hall, PO Box 7609, 2721 Founders Drive, Raleigh, NC, 27607, United States

<sup>5</sup>American Association for the Advancement of Science, Science and Technology Policy Fellow at the United States Department of Agriculture, Animal and Plant Health Inspection Service, 4700 River Road, Riverdale, MD 20737, United States

<sup>6</sup>Department of Agronomy and Plant Genetics, University of Minnesota, 1991 Buford Circle, St. Paul, MN 55108, United States

<sup>7</sup>Department of Energy, Joint Genome Institute, 2800 Mitchell Drive, Walnut Creek, CA 94598, United States

<sup>8</sup>Hudson Alpha Institute for Biotechnology, 601 Genome Way, Huntsville, AL 35806, United States

\*Corresponding author: [Steve.Larson@usda.gov](mailto:Steve.Larson@usda.gov) (tel. 01 4357971703)

**ORCID list:**

Steve Larson (0000-0003-2742-2134), Lee DeHaan (0000-0002-6368-5241), Jesse Poland (0000-0002-7856-1399), Xiaofei Zhang (0000-0003-4516-9179), Kevin Dorn (0000-0003-2722-6221), Traci Kantarski (0000-0002-8584-2965), James Anderson (0000-0003-4655-6517), Jeremy Schmutz (0000-0001-8062-9172), Jane Grimwood (0000-0002-8356-8325), Jerry Jenkins (0000-0002-7943-3997), Shengqiang Shu (0000-0002-4336-8994), Jared Crain (0000-0001-9484-8325), Matthew Robbins (0000-0002-5467-4452), Kevin Jensen (0000-0001-5560-0394)

## Abstract

Allohexaploid ( $2n=6x=42$ ) intermediate wheatgrass (*Thinopyrum intermedium*), abbreviated IWG, is an outcrossing perennial grass belonging to the tertiary gene pool of wheat. Perenniality would be a valuable option for grain production, but attempts to introgress this complex trait from wheat-*Thinopyrum* hybrids have not been commercially successful. Efforts to breed IWG itself as a dual-purpose forage and grain crop have demonstrated useful progress and applications, but grain yields are significantly less than wheat. Therefore, genetic and physical maps have been developed to accelerate domestication of IWG. Herein, these maps were used to identify quantitative trait loci (QTLs) and candidate genes associated with IWG grain production traits in a family of 266 full-sib progenies derived from two heterozygous parents, M26 and M35. Transgressive segregation was observed for 17 traits related to seed size, shattering, threshing, inflorescence capacity, fertility, stem size, and flowering time. A total of 111 QTLs were detected in 36 different regions using 3,826 genotype-by-sequence (GBS) markers in 21 linkage groups. The most prominent QTL had a LOD score of 15 with synergistic effects of 29% and 22% over the family means for seed retention and percent naked seed, respectively. Many QTLs aligned to one or more IWG gene models corresponding to 42 possible domestication orthogenes including the wheat *Q* and *RHT* genes. A cluster of seed-size and fertility QTLs showed possible alignment to a putative *Z* self-incompatibility gene, which could have detrimental grain-yield effects when genetic variability is low. These findings elucidate pathways and possible hurdles in the domestication of IWG.

## Keywords:

domestication, genetic mapping, genome sequence, perennial grains, QTL, *Thinopyrum intermedium*,

## Introduction

Intermediate wheatgrass (*Thinopyrum intermedium*) is a cool-season perennial Triticeae grass, native to parts of Eastern Europe and western Asia, that has been widely used for soil conservation and forage production in North America and other temperate regions of the World (Jensen et al. 2016; Zair et al. 2018). Intermediate wheatgrass (IWG) ranks among the highest-yielding forage and biomass crops that can be grown across regions of the Upper Midwest, Great Plains and Intermountain regions of Canada, the United States, and other places where it is cultivated (Harmony 2015; Kenneth and Kevin 2001; Larson et al. 2017; Lee et al. 2009; Monono et al. 2013; Pearson et al. 2015; Robins 2010; Wang et al. 2014). At least 15 IWG cultivars have been released for purposes of soil conservation and forage production in Canada and the United States, with some of the first cultivars selected directly from plant introductions from Russia in 1932 (Jensen et al. 2016; Knowles 1977; Pearson et al. 2015). However, selection for better fertility and seed yields was critical for development of the first widely successful North American cultivar ‘Oahe’ (Knowles 1977; Ross 1963), which is still grown today. Modern IWG forage cultivars have also undergone selection for disease resistance, forage quality, forage yield and other traits (Jensen et al. 2016; Krupinsky and Berdahl 2000; Vogel et al. 2005).

Species of the genus *Thinopyrum*, including IWG, are considered the closest perennial relatives of common wheat (*Triticum aestivum*) and comprise a useful part of its tertiary gene pool (Ceoloni et al. 2015; Lang et al. 2018; Li et al. 2017; Liu et al. 2017; Uzma et al. 2015; Zair et al. 2018). Common wheat and IWG have similar allohexaploid genomes ( $2n=6x=42$ ). However, unlike wheat, IWG is usually self-incompatible meaning that most populations are highly heterogeneous and most individuals are highly heterozygous (Jensen et al. 2016; Jensen et al. 1990; Kantarski et al. 2017; Zhang et al. 2016). Wheat-*Thinopyrum* hybrids have also been utilized for the development of perennial grain crops (Curwen-McAdams and Jones 2017; Hayes et al. 2018), but commercial production of these plant materials has not yet been realized because it has been difficult to introgress and stabilize this complex trait in wheat. However, parallel efforts to directly domesticate IWG itself as a perennial grain crop (Cox et al. 2006; Cox et al. 2010; Wagoner 1990) have led to small-scale production and utilization of IWG grain or flour for baking, beverages, and other edible food products (DeHaan and Ismail 2017). Although current grain yields are modest, IWG has potential to be used as a multipurpose perennial forage and grain crop (Bell et al. 2015; Cattani and Asselin 2017; Jungers et al. 2017; Ryan et al. 2018). These burgeoning efforts to domesticate IWG (Cattani 2017; DeHaan et al. 2018; Zhang et al. 2016) aim to diversify and enhance the quantity and quality of food products (DeHaan and Ismail 2017; Marti et al. 2016) while providing improved soil conservation, water quality, carbon sequestration, nutrient management, and other ecosystem services (Culman et al. 2013). These goals have been described as the “ecological intensification of agriculture” (DeHaan et al. 2018).

A high-density genotype-by-sequencing (GBS) consensus linkage map (Kantarski et al. 2017) and draft genome sequence were developed to accelerate the domestication of IWG (DeHaan et al. 2018). The IWG consensus map had three homoeologous sets of seven linkage groups (LG01 to LG21) with 10,029 GBS markers showing colinear alignments to the seven chromosomes of diploid barley (Kantarski et al. 2017). The consensus map was constructed using seven full-sib populations derived from 13 heterozygous individuals from the third (C3) and fourth (C4) cycle of selection (DeHaan et al. 2018) including one population derived by self-pollination of one

outstanding C3 individual, C3\_3471; a biparental population derived from two C4 parents, M26 and M35, descended from C3\_3471 and two other C3 grandparents; and another biparental population derived from two C4 parents, C4\_2856 and C4\_5353, also descended from C3\_3471 and three other C3 grandparents (Kantarski et al. 2017). The C3\_3471 individual was identified in the third cycle of selection as the first predominantly free-threshing and non-shattering IWG plant, which also had exceptionally long and heavy seeds. The IWG draft genome sequence was developed by sequencing a haploid twin-seedling (Namikawa and Kawakami 1934) from C4\_5353, which was also a parent of two mapping populations (Kantarski et al. 2017). The draft sequence includes 21 recognized chromosome sequences ranging in size from 250.8 to 802.6 megabases (MB), one unrecognized chromosome of 849.9 MB, and 1,977 scaffolds ranging in size from 0.1 to 8.8 MB with a total of 159,905 annotated gene models. The 21 chromosome sequences were numbered CHR01 to CHR21 according to alignments with the 21 LGs of the IWG GBS consensus map (Kantarski et al. 2017). The IWG consensus map has been used to identify QTLs and markers associated with seed size in two biparental populations, M26 x M35 and C3-2331 x C3-2595, and one association mapping (AM) population (Zhang et al. 2017). However, the M26 x M35 population was developed mainly to investigate the genetic control of the non-shattering and free-threshing seed traits of C3\_3471.

The inflorescence of IWG is fundamentally like wheat in that they both have one sessile spikelet at each rachis node and multiple florets per spikelet with up to one seed per floret and variable levels of fertility. Seed disarticulation in wild IWG plants normally occurs in two different ways: 1) abscission above the junction of the rachis and spikelet base producing a complex diaspore with a relatively long wedge-shaped rachis internode extending away from the spikelet similar to that produced by the brittle rachis of barley (Pourkheirandish et al. 2015) and certain types of wheat that produce wedge-shaped diaspores (Li and Gill 2006), and 2) abscission below the junction of the rachilla and floret base, within the spikelet, producing a less complex diaspore that includes a short rachilla internode extending up from the base of the floret adjacent to the palea. Threshing of semi-domesticated IWG forage varieties normally produces seeds with the lemma and palea attached to the pericarp, like those of hulled barleys. However, free-threshing forms of IWG, such as C3\_3471, and other domesticated forms of wheat and barley produce naked grains comprised of the caryopsis detached from the lemma, palea, and spikelet glumes. At least two different genes have been identified, the barley *nud* gene (Taketa et al. 2008) and wheat *Q* gene (Simons et al. 2006), which are responsible for the free-threshing trait of naked barleys and common wheat, respectively.

The domestication and improvement of IWG as a perennial grain crop currently focuses on increasing seed size, fertility, inflorescence capacity, stem and inflorescence compactness, seed retention in the field, percentage of naked seeds after threshing, and uniformity of maturity (DeHaan et al. 2018; DeHaan et al. 2016; Zhang et al. 2016). One of the primary objectives of the study herein was to identify and map QTLs controlling seed shattering, percentage of naked seeds after threshing and other traits related to reproductive fertility, inflorescence capacity, stem and inflorescence compactness, seed retention in the field, and uniformity of maturity in the M26 x M35 population using the GBS consensus map (Kantarski et al. 2017). Another major objective of this study was to compare M26 x M35 QTLs, C3-2331 x C3-2595 QTLs, and AM markers to the annotated IWG draft genome sequence and identify possible candidate genes (CGs) controlling relevant domestication and improvement traits

(Doebley et al. 2006; Kovach et al. 2007; Lenser and Theißen 2013; Meyer and Purugganan 2013; Tang et al. 2010) such as the percentage of free-threshing naked seeds (Simons et al. 2006; Taketa et al. 2008), seed shattering (Doust et al. 2014; Li and Gill 2006; Pourkheirandish et al. 2015), and grain yield (Nadolska-Orczyk et al. 2017) traits in wheat, barley, rice, and other grain crops.

## **Materials and Methods**

### **Plant materials and Field Evaluations**

The terms “parent”, “hybrid”, or “progeny” are used to describe genetically unique individuals, generally referred to as “genets”, depending on context, whereas the terms “propagule” or “plot” may refer to different clonal replicates of the same genet. A total of 266 full-sib progenies, two C4 parents (M26 and M35), and two known C3 grandparents (C3\_3471 and C3\_3941) comprising one of seven families used to construct a high-density linkage map (Kantarski et al. 2017) were clonally propagated into replicated field plots in Kansas (KS) and Utah (UT). Propagules were planted in grids with 1 m spacing in UT (1.0 m<sup>2</sup> plots) and 3 ft spacing in KS (0.81 m<sup>2</sup> plots). Although these propagules eventually grow large enough to cover most plots, tillage between propagules was used to control weeds and help maintain plot integrity. The C4 M26 genet was the maternal seed parent for 128 progenies, whereas the C4 M35 genet was the maternal seed parent for the other 138 progenies. The M26 genet originates from a cross of C3\_3471 as the seed parent and C3\_3941 as the pollen parent in the third cycle of selection for grain production traits (DeHaan et al. 2018; Zhang et al. 2016). The M35 genet originates from a cross of C3\_3941 as the seed parent and an unknown C3 pollen parent. Two propagules from each of 221 progenies, two C4 parents, and two known C3 grandparents were transplanted at the KS location (38.771517° N / -97.569408° W) on November 28, 2012. Three clones from each of 253 progenies, two parents, and two known grandparents were transplanted into a field in UT (41.695957° N / -111.831358° W) on May 6, 2013.

A total of seventeen seed production traits were measured on each plot once per harvest year for at least two years (2014 and 2015) at both locations (KS and UT) and most traits were also evaluated at the KS location in 2013. The KS location was planted early enough in 2012 to enable sufficient plant growth, vernalization, and flowering to evaluate seed production traits in 2013. Conversely, the UT location was planted in the spring of 2013, which did not allow sufficient growth, vernalization, flowering, and set seed to evaluate seed production traits in 2013.

Four traits measured in the field include crown circumference (CRCI), number of inflorescences per crown (INCR), stem length (STLE), and one Zadoks (Zadoks et al. 1974) maturity rating (ZAMA) on each plot, once per year. The CRCI and INCR traits were evaluated by measuring the distance (cm) around the outermost stems at the base of each propagule, just above the soil surface, and then counting the number of inflorescences in each plot in 2014 at both locations. These traits were not evaluated in 2015 in KS because it was too difficult to discern the then natural edges of different propagules. However, in UT, propagules were just beginning to grow into cultivated spaces between rows in 2015, so we devised a faster and possibly more effective way to estimate CRCI and INCR. First, we counted the number of stems through a 10-cm wide section of the widest undisturbed portion of each crown using an open-ended rectangular quadrat with marked distances along each arm to measure the maximum

diameter ( $d$ ) of the crown. Estimates of CRCI, from UT in 2015, were then determined based on the formula for the circumference of a circle ( $C = \pi \cdot d$ ). Estimates of INCR, from UT in 2015, were then determined using the formula for the area of a circular crown [ $A = \pi \cdot (0.5 \cdot d)^2$ ] multiplied by the tiller density [ $S/(10 \cdot d)$ ]. The total (stretched) STLE was measured (cm) from the soil surface to the upper tip of the inflorescence. The peduncle and inflorescence from ten of the tallest culms were harvested from each plot for subsequent measurements of 13 seed and inflorescence traits in the laboratory.

Five stem and inflorescence traits including the stem width (STWI), number of spikelets per inflorescence (SPIN), inflorescence length (INLE), seed shattering (SESH), and number florets per spikelet (FLSP) were measured on a subset of three of 10 harvested culms from each plot. The INLE was measured from the rachis node subtending the lowest spikelet to uppermost point of the inflorescence. The SESH at the KS location was determined based on the percentage of disarticulation after bending spikelets to 90° angle from the rachis and dropping inflorescences from a height of 25 cm. The SESH at the UT location was determined by the percentage of disarticulation after three repeated strikes of each stem to a table surface. The STWI was measured on the thinnest part of each stem just below the inflorescence.

For UT field evaluations, six seed traits were measured after removing seed from ten harvested culms using a LD 180 laboratory thresher (Wintersteiger Inc., Salt Lake City, Utah, USA). Seeds and chaff were further separated using a General Seed Blower (Seedburo Equipment Co., Des Plaines, Illinois, USA). The percentage of seeds threshed out naked (SENA), seed area (SEAR), seed length (SELE), seed width (SEWI), and total number of seeds were determined using SmartGrain phenotyping software (Tanabata et al. 2012) from images of cleaned seeds from all ten inflorescences taken from a scanning device with a blue-paper background. The SENA was determined by first counting the number of hulled seeds and then counting the number naked seeds based on color recognition of lighter colored hulls and darker colored pericarps, respectively. Measurements of SEAR, SELE, and SEWI were also taken from the darker colored naked seeds, excluding broken parts less than 2.0 mm long clumps of seeds that exceeded 8.0 mm length based on visual validation of SmartGrain image annotations. The total seed yield per inflorescence (SYIN) and average seed mass (SEMA) were determined by dividing the mass of all cleaned seeds (mg) by the number of inflorescences harvested (10) and the total number of seeds, respectively.

For KS field evaluations, six seed traits were measured after removing seed from ten harvested culms using a spike-tooth small bundle thresher. Additional cleaning was performed by hand sieving with a 12/64 inch round hole sieve and aspirating with an STS-WM2 Air Separator (Seed Tech Systems, Wilton, CA, USA). The percentage of seed threshed out naked (SENA) was visually estimate for each sample. Approximately 20 naked seeds from each sample were photographed and the images were analyzed in ImageJ (Schneider et al. 2012). Image analysis provided seed area (SEAR), seed length (SELE), seed width (SEWI), and number of seeds. This subsample was weighed and the seed number data was used to calculate average seed mass (SEMA). The total seed yield per inflorescence (SYIN) was determined by dividing the mass of all cleaned seeds by the number of inflorescences harvested (10).

Two other seed- and inflorescence-related traits, the number of seeds per spikelet (SESP) and the number of seeds per floret (SEFL), were calculated based on measurements of the average number of seeds per inflorescence, SPIN, and FLSP as described above.

## **Data Analyses**

Least square means (LSMEANS) for each genet and least significant differences (LSD) among genets were determined using the SAS version 9.4 (SAS Institute, Cary, North Carolina, USA) MIXED procedure with genets, years, locations, and the three-way interaction term as fixed effects and replications as random effects. The overall means and LSD categories ( $p \leq 0.05$ ) for the grandparents and parents were determined using the repeated option to model covariance structure between years. Segregation of the trait values used for QTL analysis of the full-sib progeny, from LSMEANS procedure, was measured in part by relative standard deviations (standard deviation among trait values / mean over all trait values) which reflects the magnitude of trait variability, relative to the overall mean, among 68% of the most typical individuals in a normally distributed population. The broad-sense heritability ( $H$ ) estimates and standard errors were determined on a single plot basis and entry (genet) mean basis for randomized complete block designs in multiple environments using a SAS MIXED procedure with environments (five location x year combinations), replications within environments, genet, and genet x environment as random effects (Holland et al. 2010). Pearson correlation tests were performed for 136 pairwise comparisons of the 17 traits, based on individual plot measurements, using environments (locations and years) as a grouping factor for the statsBy function of the R (R Core Team 2017) psych package (Revelle 2018). Significance thresholds for the trait correlation tests were adjusted using a Bonferroni correction to control for multiple testing.

Two different approaches for QTL detection based on models for a two-way pseudo-testcross (TWPT) and cross-pollinated (CP) plants, more fully described in following paragraphs, were performed using MapQTL version 6 (Van Ooijen 2009). All of the map files and locus data used for these QTL analyses were based on the integrated GBS consensus map (Kantarski et al. 2017) with a total of 3156 markers from the M26xM36 family. This map included 1699 markers that were heterozygous in M26 (progeny genotypes *lm* or *ll*), 1087 markers that were heterozygous in M35 (progeny genotypes *nn* or *nm*), and another 1070 markers there were heterozygous in both M26 and M35 (progeny genotypes *hh*, *hk*, or *kk*). The quantitative trait data were based on LSMEANS trait estimates for each genet, within and among environments, as described above. All TWPT and CP QTL analysis were initially performed using the same single-QTL interval mapping procedure to identify possible QTL markers that were subsequently used as cofactors in the first round of restricted multiple-QTL model (rMQM) mapping (Van Ooijen 2009). A second round of rMQM mapping was also performed for all TWPT and CP QTL analyses using cofactors for QTLs that were significant after the first round of rMQM mapping. All QTL analyses were performed using a maximum likelihood mixture model, with up to 20 iterations, to determine the LOD likelihood ratio statistic (Van Ooijen 2009). A permutation test for each CP and TWPT analysis was used to determine significance thresholds for the LOD statistic corresponding to a genome-wide P-value of 0.05 (5%) to identify putative QTLs and cofactors (Van Ooijen 2009). Only those QTLs exceeding the 5% genome-wide LOD threshold in the second round of rMQM mapping were considered significant.



The two-way pseudo-testcross (TWPT) QTL analysis utilized a doubled haploid model with separate maps for each parent (Van Ooijen 2009), M26 and M35. Briefly, the *lm* and *ll* genotypes from 1699 M26 markers were changed to *A* and *B*, respectively; and the *nm* and *np* genotypes from 1087 M35 markers were changed to *B* and *A*, respectively. The remaining 370 markers with genotypes *hh*, *hk*, or *kk* were deleted. A map with 42 LGs including 1699 M26 markers in 21 LGs and 1087 M35 makers in another 21 LGs was assembled using marker positions from the integrated consensus map (Kantarski et al. 2017). Only one genotypic effect, the difference between *A* or *B* marker alleles from one parent, is fitted at any given map position in the TWPT approach. The TWPT design is a simplified model that has possible theoretical advantages because it enables rMQM mapping of each LG from each parent using QTL marker cofactors from other LGs of both parents, including one homologous LG from the other parent (Van Ooijen 2009). Another practical advantage of the TWPT design is that it is relatively easy to identify useful QTL cofactors when the parental LGs are separated since some QTLs may not be heterozygous both parents. Thus, to identify TWPT rMQM cofactors we initially selected one marker with the highest LOD score overall environments for each M26 or M35 LG having at least one significant TWTP QTL. However, the automatic cofactor selection procedure of MapQTL 6 was also used to identify the final set of TWPT rMQM cofactors.

The second approach for QTL analysis utilized an integrated map containing 3156 markers in 21 LGs for both M26 and M35 parents (Kantarski et al. 2017), which was constructed using the cross-pollinator (CP) model (Van Ooijen 2006). In full-sib CP families, one or more QTLs may be heterozygous in one or both parents with up to four possible alleles per QTL. The CP QTL approach always fits four possible QTL alleles designated *a* and *b* corresponding to marker alleles *l* and *m*, respectively, of the first parent (M26) and QTL alleles *c* and *d* corresponding to marker alleles *n* and *p*, respectively, of the second parent (M35). The CP QTL analysis (Van Ooijen 2009) has theoretical and practical advantages in that three possible genotypic effects are fitted including the difference between *a* and *b* QTL alleles of the first parent ( $\alpha$ ), the difference between *c* and *d* QTL alleles of the second parent ( $\gamma$ ), and the intralocus interaction ( $\tau$ ) as deviations from the overall mean ( $\mu$ ). Thus, in a cross of *ab* x *cd* QTL alleles the expected progeny phenotypes are modeled (Van Ooijen 2009) as follows:

$$ac = \mu - \alpha - \gamma - \tau$$

$$ad = \mu - \alpha + \gamma + \tau$$

$$bc = \mu + \alpha - \gamma + \tau$$

$$bd = \mu + \alpha + \gamma - \tau$$

If the parents (M26 and M35) are heterozygous for the same two QTL alleles, *a* and *b*, then  $\tau$  would represent a dominance deviation term. However, this is never assumed to be the case because MapQTL 6 CP model always fits separate effects,  $\alpha$  and  $\gamma$ , for both parents. Since more than 72% of the markers in the integrated CP map were not informative in one parent or the other parent (Kantarski et al. 2017), the interval mapping procedure can assign relatively high LOD scores to M26 or M35 marker loci that may not be directly associated with the QTL. Thus, in addition to the TWPT rMQM cofactors, the Kruskal-Wallis procedure of MapQTL 6 was used to help identify additional CP rMQM cofactors for each CP QTL. However, we did not consider more than one informative marker per parent per LG for use as a possible CP rMQM cofactor.

## Comparative mapping

The two-LOD drop off intervals for M26 x M35 QTLs with the highest LOD values for each trait and each LG, including parent-specific TWPT QTLs, were graphed onto the integrated M26 x M35 GBS consensus map (Kantarski et al. 2017) using the R (R Core Team 2017) LinkageMapView package (Ouellette et al. 2018). Another 44 SEMA, SEAR, SELE, and SEWI QTLs, detected over one or more years in the C3-2331 x C3-2595 (UMN) family (Zhang et al. 2017), were included on this graph for comparison to M26 x M35 QTLs detected in this study. The probability ( $P_{xy}$ ) that one of the  $x$  most significant QTL markers was the same as one of the  $y$  most significant association-mapping markers, with a total of  $s$  shared markers, was calculated as one minus the probability of not having any of the same markers, which equals the products of one minus the chance of drawing one of  $y$  numbers  $x$  times (where the total number of possible makers,  $s$ , decreases from  $i = 0$  to  $x-1$  without replacement) or drawing one of  $x$  numbers  $y$  times (where the total number of possible makers,  $s$ , decreases from  $i = 0$  to  $y-1$  without replacement) as follows:

$$1 - \left( \prod_{i=0}^{x-1} \left( 1 - \left( \frac{y}{s-i} \right) \right) \right) \text{ or } \prod_{i=0}^{y-1} \left( 1 - \left( \frac{x}{s-i} \right) \right) = P_{xy}.$$

Sequences of GBS markers (Kantarski et al. 2017), including 51 significant AM markers (Zhang et al. 2017), were aligned to the pre-publication “*Thinopyrum intermedium* C4-5353-T1 Annotated Standard Draft” sequence, available on the JGI Genome Portal (<https://genome.jgi.doe.gov/portal>), using a Basic Linear Alignment Search Tool (BLAST) with an Expect-value ( $E$ ) threshold  $\leq 1e-10$  (Altschul et al. 1990). Graphical comparisons of the 21 LGs (LG01 to LG21) and 21 corresponding chromosome sequences (CHR01 to CHR21), based on GBS marker alignments, were performed using LinkageMapView by normalizing LG lengths, measured in centiMorgans (cM), and CHR lengths, measured in nucleotide mega-bases (MB).

Seed-yield and domestication orthogenes of wheat, barley, rice, maize, and other plants (Doebley et al. 2006; Doust et al. 2014; Hackauf and Wehling 2005; Lenser and Theißen 2013; Li and Gill 2006; Meyer and Purugganan 2013; Nadolska-Orczyk et al. 2017; Pourkheirandish et al. 2015; Simons et al. 2006; Taketa et al. 2008; Tang et al. 2010) including *Arabidopsis* (Balanza et al. 2016) and *Lolium* (Manzanares et al. 2016; Shinozuka et al. 2010) were aligned to the IWG draft genome sequence using BLAST or BLASTX (Altschul et al. 1997) with a minimum significance threshold of  $E \leq 1e-20$ . At least three BLAST or BLASTX hits corresponding to three possible orthogenes on three homoeologous chromosomes of IWG (Kantarski et al. 2017) were considered for each CG. If the three most significant IWG BLAST hits were located on chromosome sequences with known orthology to chromosomes harboring barley, *Lolium*, rice, or wheat CGs (Klaas et al. 2011; La Rota and Sorrells 2004; Thorogood et al. 2017; Tulpan and Leger 2017) then only these hits were considered as possible orthogenes. Additional IWG BLAST hits having similar  $E$  values were considered when queried using *Arabidopsis* genes allowing for ancient duplications and many possible genome rearrangements that have occurred between dicots, such as *Arabidopsis*, and monocots such as IWG. Only those IWG BLAST hits corresponding to annotated IWG gene models, supported by IWG transcripts, were considered as possible IWG CGs.

## Results

### Phenotypic and genotypic variation

Significant differences ( $p \leq 0.05$ ) were detected among the C3 grandparents and C4 parents for all 17 traits, with up to four levels of difference for SEWI, SYIN, and ZAMA (Table 1). Moreover, significant effects ( $p \leq 0.05$ ) of genet, year, location, and three-way interaction term were detected for all traits except FLSP, SEFL and INLE in the overall analysis of the grandparents, parents, and 266 full-sib progenies. In most cases, those effects were highly significant ( $p \leq 0.0001$ ), therefore all further analyses were conducted for each location x year environment. Only the year effect was not significant for INLE and only the location effect was not significant for FLSP and SEFL. The standard deviations among the progeny trait estimates were within 4% to 8% of the overall trait means for SELE, SEWI, and ZAMA (Table 2). Conversely, the standard deviations varied from 38% to 86% of the trait means for SEFL, SESH, SESP, and SYIN (Table 2). However, broad-sense heritability ( $H$ ) estimates for traits with lower variability tended to be greater than  $H$  estimates for traits with high variability (Table 2). The relative standard deviations for SENA varied from 17% to 94% among environments, in part because the overall means of this trait also varied widely among environments (Table 2). These data, Tables 1 and 2, demonstrate significant genetic variation among the C3 grandparents, C4 parents, and full-sib progeny, respectively, for all 17 traits.

Significant correlations ( $p < 0.05$ ) were detected for 100 of the 136 possible pairwise comparisons among 17 traits, controlling for multiple testing (Table 3). Although there were up to 12 possible plot measurements taken for each of the 270 genets (two replications in KS evaluated three years and three replications in UT evaluated over two years), with 3240 possible measurements per trait, the maximum number of observations was limited to 2570 measurements because not all genets were present in all five replications. Relatively strong positive correlations were observed among the four seed-size traits (SEAR, SELE, SEMA, and SEWI) and between the two fertility traits, SEFL and SESP (Table 3). Seed yield per inflorescence (SYIN) is a complex trait that showed relatively strong and significant correlations with fertility traits, SEFL and SESP, and moderate correlations with seed size (SEAR, SELE, SEMA, and SEWI), inflorescence capacity (SPIN and FLSP), and seed disarticulation (SENA and SESH) traits (Table 3). The percentage of naked seeds (SENA) was negatively correlated SEWI, SESH, and FLSP (Table 3), which was especially true in the KS evaluations where these correlation coefficients were -0.25, -0.34, and -0.12, respectively.

### QTL analysis

Permutation tests showed that the minimum LOD thresholds required to control for a 5% genome-wide error rate ( $P < 0.05$ ) were 3.5 and 4.7 or less for the TWPT and CP QTL analyses, respectively, for each trait. Using these LOD thresholds for all 17 traits, there were a total of 210 significant QTLs detected among six QTL analyses for each of five different environments and averages over all five environments (Table 4; Supplemental Document 1). A subset of 56 QTLs were significant in only one analysis, whereas 55 QTLs were significant among two or more analyses (Supplemental Document 1). A total of 57 QTLs were significant in the analysis of trait averages over all five environments but only one QTL, for SELE on LG06, was significant across all six analyses (Table 4). Considering the most significant QTL with maximum LOD on each LG and each trait, there were at least 111 distinct QTLs

including 19 M26 and 20 M35 QTLs detected using the TWPT model and 72 QTLs detected using the CP model (Tables 4 and 5; Supplemental Document 1). Only 3 distinct QTLs were detected for CRCI and up to 10 distinct QTLs were detected for ZAMA (Table 5). There were at least two significant QTLs on each of the 21 LGs, with up to 12 significant QTLs on LG06 (Table 5).

For most traits except INLE, the total magnitudes of maximum QTL effects (Max-total) and overall average QTL effects (Avg-total), within and among five environments, were not the same for both M26 and M35 parents (Table 6). The Max-total and Avg-total M35- $\gamma$  effects were greater than corresponding M26- $\alpha$  effects for SEMA, FLSP, SEFL, SESP, STWI, ZAMA, CIRCI, and INCR (Table 6). Conversely, the Max-total and Avg-total M26- $\alpha$  effects were greater than corresponding M35- $\gamma$  effects for SEAR, SENA, SYIN, and STLE (Table 6). The Avg-total M26- $\alpha$  effects were also greater than corresponding M35- $\gamma$  effects for SELE, SESH and SPIN, but this was not true for the Max-total effects of these four traits (Table 6). A more detailed examination of SENA QTL effects indicates the magnitude of M26- $\alpha$  QTL effects were equal or greater than the M35- $\gamma$  effects for six of nine SENA QTLs including the LG02 and LG11 SENA QTLs, which had relatively large M26- $\alpha$  effects (Table 6). Thus, the Avg-total M26- $\alpha$  SENA effects (28% of population mean) was about 133% greater corresponding M35- $\gamma$  effects (12% of population mean), with a total estimated effect of 52% of the SENA mean (Table 6). The LG02 SENA QTL was most significant in the KS15 analysis (Table 4), where progenies with LG02 SENA QTL genotypes *ac* and *bd* were 39% greater than the population mean or 25% less than the population mean, respectively (Table 6; Supplemental Document 1). If the LG02 M26 *a* and *b* alleles are additive, then it could be inferred that the SENA means for progeny with an *aa* genotype would be 42% over the population mean (Table 2), with a M26- $\alpha$  SENA effect of 21% of population mean (Table 6). However, there are no progeny or parents that are homozygous for the four possible alleles (*a*, *b*, *c*, or *d*) of any QTLs in the M26 x M35 family so it is not possible to determine the effects possible QTL genotypes. Nevertheless, the Avg-total SENA QTL effects, 68% of population mean (Table 6), were enough to account for the range of 23.2% to 83.5% SENA variation (55.3% average) among progeny (Table 2). Moreover, the Avg-total M26- $\alpha$  effects, 28% of population mean (Table 6), would also be enough to account for differences of 28.7% and 94.1% SENA between the parents of M26, C3\_3941 and C3\_3471 (Table 1), if they were homozygous for small- and large-effect alleles, respectively, and if the M26 *a* and *b* alleles are additive.

The highest LOD score for all 17 traits was for the LG11 SESH QTL (Table 5), which had relatively strong but opposite effects on both SESH and SENA (Table 6) meaning that it has a desirable or synergistic effects of decreasing seed shattering and increasing the percentage of free-threshing (naked) seed. In fact, this LG11 QTL was the only QTL that had synergistic M26- $\alpha$  SESH and SENA QTL effects and synergistic M35- $\gamma$  SESH and SENA QTL effects (Table 6). Only one other QTL, on LG10, had synergistic SESH and SENA M26- $\alpha$  effects (Table 6). The total magnitude of the LG10 and LG11 SESH and SENA M26- $\alpha$  effects, 23% and 26% respectively (Table 6), would be enough to account for most SESH and SENA differences between the parents of M26, C3\_3941 and C3\_3471 (Table 1), if they were homozygous for desirable and undesirable alleles, respectively, and if the M26 *a* and *b* alleles are additive. Two other QTLs on LG12 and LG14 had synergistic SESH and SENA M35- $\gamma$  effects. Thus, at four QTLs on LG10, LG11, LG12, and LG14 had synergistic M26- $\alpha$  or M35- $\gamma$  effects on SESH and SENA (Table 6), which account for negative correlations between these traits (Table 3). Only one QTL, on LG06, had the

same positive or negative effect on SESH and SENA (Table 6). In general, the relative directions of QTL effects for different traits (Table 6) were consistent with trait correlations (Table 3). For example, nine of the 13 LGs associated with seed size traits (SEAR, SELE, SEMA, and SEWI) had the same directional effects for  $\alpha$  or  $\gamma$  on more than one of these traits, which was consistent with positive correlations among these traits (Table 3). Likewise, six of the 10 LGs associated with SEFL and SESP had the same directional  $\alpha$  or  $\gamma$  effects for both traits, consistent with the positive correlation between these two traits (Table 3).

The percent variation explained (PVE) by M26 was substantially greater than M35 for SEAR, SELE, SENA, SPIN, FLSP, and STLE (Table 7). Conversely, the PVE by M35 was substantially greater than M26 for SEMA, SEWI, SEFL, SESP, INLE, STWI, ZAMA, CRCI, and INCR (Table 7). Thus, the relative PVE by M26 and M35 (Table 7) closely mirrors the relative magnitudes of the average M26- $\alpha$  and M35- $\gamma$  QTL effects among all five environments (Table 6). However, the combined PVE by M26 and M35 in the CP approach was greater than the combined PVE by M26 and M35 in the TWPT approach for SEMA, SEWI, SPIN, ZAMA, and CRCI (Table 7), presumably because the CP model includes effects of intralocus interactions ( $\tau$ ).

#### **Alignment of IWG QTL intervals and association mapping (AM) loci to IWG draft genome sequence**

A subset of 3,608 (93.6%) of the 3,856 GBS markers comprising the M26 x M26 linkage map had significant matches to the IWG physical map. Of these 3,608 matches, 2,977 (82.5%) showed synteny with homologous chromosome sequences such as LG01 and CHR01 (Figure 1), 239 (6.6%) mapped to different members of the same homoeologous group (HG) such as LG01 and CHR02 or CHR03, 213 (5.8%) mapped to different genetic and physical HGs, and 179 (5.0%) matched unmapped scaffolds. Substantial collinearity among GBS markers of the 21 linkage groups and corresponding chromosome sequences was also discernable (Figure 1). A subset of 1,072 (29.7%) of the 3,608 M26 x M35 GBS marker alignments were located within 1,072 (0.7%) of the 159,905 IWG gene models. A slightly higher portion of 37 (33%) of the 111 most significant M26 x M35 QTL markers were located within IWG gene models. A total of 9,403 (93.7%) of the 10,029 GBS markers from the IWG consensus map (Kantarski et al. 2017) showed significant matches to the draft genome sequence, including 3,223 (34.3%) located within annotated IWG gene models.

A total of 93 GBS markers had the highest LOD score for the 111 most significant M26 x M35 QTLs and the two-LOD drop off intervals for many of the M26 x M35 QTLs were identical or very similar (Supplemental Document 1). The 111 M26 x M35 QTLs collapsed into 72 QTL intervals spanning different regions or different lengths of the linkage map (Figure 1). However, these 72 QTL intervals covered no more than 36 non-overlapping regions of the 21 integrated LGs. One region of LG06 had significant effects on 11 different traits including all seed-size, fertility, and seed harvest traits (Figure 1).

A total of 44 SEMA, SEAR, SELE, and SEWI QTLs from the biparental UMN C3-2331 x C3-2595 population (Zhang et al. 2017) were mapped to 22 QTL intervals corresponding to 21 non-overlapping regions of the IWG consensus map (Figure 1). Similarly, 25 M26 x M35 seed-size QTLs were mapped to 20 QTL intervals corresponding to only 14 non-overlapping regions of the same IWG consensus map (Figure 1). A total of 69 seed QTLs from both families, M26 x M35 and C3-2331 x C3-2595, mapped to only 26 non-overlapping regions on 20

LGs with correspondence between families on LG06, LG08, LG09, LG11, LG14, LG15, LG17, and LG20 (Figure 1). A total of 51 AM markers mapped to 18 of the 21 LGs, excluding LG06, LG07 and LG19 (Zhang et al. 2017). No significant seed-size QTLs were detected on LG05, in the M26 x M35 or C3-2331 x C3-2595 families, but there were eight AM markers on this LG. Five LGs (LG02, LG06, LG15, LG17, and LG20) had significant QTL effects on all four seed-size traits (SEMA, SEAR, SELE, and SEWI), but only LG06 had significant QTL effects on all four seed-size traits in both M26 x M35 and C3-2331 x C3-2595 families. Only LG03, LG05, LG13 and LG20 had significant AM markers for all four seed-size traits. Only LG20 had significant QTL and AM markers for all four seed-size traits.

Two of the 25 most-significant M26 x M35 and 44 most-significant C3-2331 x C3-2595 UMN seed-size QTL markers, TP678810 and TP693406, were also among 51 most significant seed-size AM markers. The TP678810 GBS polymorphism was the most significant marker for the LG17 SELE QTL in the M26 x M35 family (Supplemental Document 1) and it was also associated with SEMA, SEAR, and SELE in the UMN AM population (Zhang et al. 2017). The TP693406 polymorphism was the most significant marker for a SEWI QTL (*Ti\_QSws.umn\_4.1*) in the C3-2331 x C3-2595 family and the SELE in the AM population (Zhang et al. 2017). The coincidence of these two markers, ranking among the 69 most significant QTL markers and 51 most significant AM markers, raised questions about the probability of this happening by chance considering that there were 3,856 M26 x M26 markers, 2,167 C3-2331 x C3-2595 markers, and 4,873 AM markers used in these experiments (Zhang et al. 2017). The probability of having at least one identical marker between the M26 x M35 QTL experiment and AM experiment was calculated as 13.9% using the formula for  $P_{xy}$  where only  $x=12$  of the most significant M26 x M35 QTL markers,  $y=14$  AM markers, and  $s=1,139$  markers were common to both experiments. The probability of having at least one identical marker in the C3-2331 x C3-2595 QTL experiment and AM experiment was calculated as 18.9% using the formula for  $P_{xy}$  where only  $x=23$  of the most significant C3-2331 x C3-2595 markers,  $y=10$  of the most significant AM markers, and  $s=1,115$  markers were common to both experiments. The probability of both occurrences was 2.6%, which suggests that these coincidences occurred more frequently than expected by chance alone. These two shared markers, LG17 TP678810 and LG04 TP693406, were located within two IWG gene models, Thintv21245331m.g and Thintv21054937m.g, respectively. The LG17 Thintv21245331m.g IWG gene model belongs to a *GDSL lipase (GLIP)* gene and the LG04 Thintv21054937m.g IWG gene model encodes a putative *xylogalacturonan beta-1,3-xylosyltransferase, xylogalacturonan deficient (XGD)*, gene (Supplemental Document 2). A total of 22 (43%) of the 51 most significant AM markers (Zhang et al. 2017) were located in annotated IWG gene models whereas only 34.3% of all 9,403 GBS alignments to the IWG draft genome sequence were located in structurally annotated gene model, suggesting that some of the most significant AM markers may be located in functionally relevant genes.

Alignments of 42 domestication genes of wheat, barley, rice, and other plants (Table 8) to the draft genome sequence of IWG detected significant homology to 142 annotated gene models of IWG (Figure 1; Supplemental Document 2). Most of these genes showed evidence of orthology based on functional annotations and known patterns of synteny among species (Supplemental Document 2). Moreover, alignments of IWG chromosome

sequences to the IWG QTL map, based on GBS markers, revealed 98 CGs with possible alignments to one or more relevant IWG QTLs (Table 8, Figure 1; Supplemental Document 2).

## Discussion

### IWG QTLs correspond to candidate genes (CGs) on seven homoeologous groups (HGs)

One of the ultimate goals of many QTL studies is to identify genes and mutations that underlie functionally important traits, initiated here by the identification of IWG loci corresponding to domestication orthogenes of closely related species such as wheat, barley, and rice. The development of an annotated draft genome sequence for IWG greatly facilitated this effort with identification of 142 possible domestication orthogenes, including 98 possible CG-QTL alignments. Discussion of these alignments was organized according to the seven homoeologous groups of wheat, barley, and IWG (Kantarski et al. 2017), which also have well-defined relationships to the 10 chromosomes of maize and 12 chromosomes of rice (La Rota and Sorrells 2004; Tulpan and Leger 2017).

Homoeologous group 1 was interesting in part because it produced the highest SENA LOD score (Table 5). Two possible orthologs of the *Seedstick* gene of *Arabidopsis* (Balanà et al. 2016), designated *STKa* and *STKb*, were found on each of the three IWG HG1 chromosomes (Figure 1). The *Seedstick* gene encodes a MADS-box transcription factor required for seed abscission in *Arabidopsis*, with seeds remaining attached to the funiculus when after fruits dehisce in *stk* mutants (Balanà et al. 2016). Although patterns of synteny between *Arabidopsis* and IWG are not easy to discern (Tulpan and Leger 2017), the functional annotation of the putative IWG *STK* orthogenes match that of *Arabidopsis* (Supplemental Document 2). The IWG *STKa* and *STKb* loci on CHR02 align to a SENA QTL on LG02, which had the most consistent SENA effects across environments (Table 4) and the highest LOD score (Table 5). Orthologs of the rice *Grain weight chromosome-5 QTL (GW5)* gene (Shomura et al. 2008; Weng et al. 2008) aligned to seed-size QTL on LG01, LG02, and LG03 (Table 8). One of three possible IWG orthologs of maize the *Barren inflorescence 4 (Bif4)* gene (Galli et al. 2015) located on CHR01, CHR02, and CHR03 aligned to a SPIN QTL on LG03. The maize *Bif1* and *Bif4* genes encode AUXIN/INDOLE-3-ACETIC ACID (Aux/IAA) proteins required for early steps of inflorescence development (Galli et al. 2015).

Homoeologous group 2, particularly LG06, produced the highest M26 x M35 LOD scores for SEMA, SEAR, SELE, and SYIN. The LG06 SELE QTL had the second highest LOD score of 12.0 in this experiment (Table 5) and it was the only QTL that was significant across all five environments (Table 6). Homoeologous group 2 was also remarkable in that it showed significant M26 x M35 QTL effects for all traits except CRCI and INCR (Table 5) and the greatest overall number of seed-size QTLs (22) in the M26 x M35 family (four QTLs), C3-2331 x C3-2595 UMN family (7 QTLs), and AM population (11). The *Grain incomplete filling (GIF)* cell-wall invertase gene of rice (Li et al. 2013; Wang et al. 2008; Yan et al. 2011), *grain length chromosome-7 QTL (GL7)* gene (Wang et al. 2015), *Sucrose synthase 2 (SUS2)* gene (Hou et al. 2014; Jiang et al. 2011), and *xylogalacturonan-deficient (XGD)* gene of *Arabidopsis* (Jensen et al. 2008) were present on CHR04, CHR05, and CHR06 with alignments to seed-size and seed yield traits on LG04 and LG06 (Figure 1, Table 8). A *GL7* locus on LG06 is an interesting candidate for the major-effect SELE QTL associated with this linkage group (Figure 1, Tables 5 and 6). The *XGD* locus on CHR04 was of particular interest because it contained the most significant SEWI QTL marker in the IWG C3-2331 x C3-2595 family, TP693406, which was also associated with SELE in the AM population (Zhang et al.

2017). The *xgd1* mutant of *Arabidopsis* has decreased levels of xylose and pectic xylogalacturonan, which are important components of cell walls and reproductive tissues that may affect process of plant development such as pectin degradation during fruit ripening (Jensen et al. 2008). The *Clustered primary branch 1 (CPBI)* gene encodes a cytochrome P450 protein involved in brassinosteroid biosynthesis, which is associated with inflorescence architecture, seed size, and plant height traits including the *DWARF11 (D11)* phenotype of rice (Wu et al. 2016). Three *CPBI* loci were found on CHR05, CHR06, and one scaffold that presumably should map to CHR04 where it might align to inflorescence and seed-size QTLs on LG04. The CHR05 and CHR06 *CPBI* genes aligned to STLE and seed-size QTLs on LG05 and LG06, respectively. Candidate genes closely associated with the barley gibberellin-insensitive *Semidwarf 3 (SDW3)* gene (Vu et al. 2010) were present on CHR04, CHR05, and CHR06 including one that also aligned to the relatively broad STLE QTL interval on LG05 (Figure 1). The *Six-rowed spike 1 (Vrs1)* encodes a homeobox-leucine zipper protein that reverts the rudimentary lateral spikelets of two-row barleys into fully developed spikelets of the six-row barleys (Komatsuda et al. 2007). Homoeologous *Vrs1* genes were present on CHR04, CHR05, and CHR06 with two copies of this gene on CHR06. The *Vrs1* genes on CHR04 and CHR06 align to SPIN and fertility (SEFL and SESP) QTLs on LG04 and LG06, respectively (Figure 1, Table 8). The barley *ZEOCRITON (ZEO)* spike density gene, located on the long arm of barley chromosome 2H (Houston et al. 2013), was present on the long arm of IWG CHR04, the short arm of IWG CHR06, and a paralogous locus on IWG CHR21 (Figure 1). The IWG ZEO loci did not align to INLE QTLs, but there was an INLE QTL aligned to the long arm of IWG CHR05, which may not be fully and correctly assembled.

Homoeologous group 3 had the highest LOD scores for SEWI, SPIN, FLSP, and INLE (Table 5). A total of 17 loci corresponding to six HG3 CGs (Table 8) were syntenous and colinear among CHR07, CHR08, and CHR09 (Figure 1). Mutations of the maize *Barren inflorescence 1 (BIF1)* gene (Galli et al. 2015) and *Barren stalk 1 (BA1)* gene (Gallavotti et al. 2004) reduce the number of spikelets due to defects in auxin signaling. The barley chromosome-3H *Six-rowed spike 4 (VRS4)* gene, an ortholog of the maize *RAMOSA2* inflorescence architecture gene, is associated with spikelet fertility and determinacy (Koppolu et al. 2013). The *BA1*, *BIF1* and *VRS4* genes aligned within or near SPIN and FLSP QTL intervals on LG07 and LG09. The *Grain number chromosome-1 QTL (GNI)* gene (Ashikari et al. 2005; Zhang et al. 2012a) is a cytokinin oxidase gene associated with a major grain-number QTL and grain weight in rice. The *GNI* gene was present on CHR07 and two CHR08 loci aligned to SEAR and SPIN QTLs on LG07 and a SEWI QTL on LG08. The *Brassinosteroid-insensitive 1 (BR1)* gene is responsible for the *uzu* semidwarfing mutation (Chono et al. 2003), which has pleiotropic effects on spike length (Chen et al. 2016) and aligned to INLE and STLE QTLs on LG07 and LG09. The barley *DENSO (Semidwarf 1)* gene encodes a gibberellic acid (GA)-20 oxidase enzyme required for GA biosynthesis (Jia et al. 2009), which can have pleiotropic effects on heading date and possibly grain size but not spike length (Kuczyńska et al. 2014). The *DENSO* gene aligned within or near STLE and seed-size QTLs on LG08 and LG09. The rice *Grain size chromosome-5 QTL (GS5)* serine carboxypeptidase gene plays a major role in regulating grain size and weight in rice and possibly wheat (Li et al. 2011; Ma et al. 2016). The *GS5* gene maps to wheat 3A, 3B, and 3D (Ma et al. 2016) and IWG CHR07, CHR08, and CHR09 (Figure 1), which align to seed size QTLs on LG07, LG08, and LG09.



Homoeologous group 4 produced a total of 23 M26 x M35 QTLs, only slightly fewer than HG5, and produced the highest LOD scores for STLE, SESH, ZAMA, and INCR (Table 5). This HG was exceptional in that it produced the highest LOD score for any trait, which was associated with the LG11 SESH QTL (Table 5). Moreover, significant QTL effects for SESH, SENA, SEFL, and ZAMA were detected on all three HG4 linkage groups (LG10, LG11, and LG12). Orthologs of the wheat ‘green revolution’ *Reduced height (RHT)* gene (Peng et al. 1999) were found on CHR10 and CHR12 including one that aligns to the high-LOD STLE QTL on LG10, which also overlaps with other QTLs controlling other traits (Figure 9). Despite efforts to find seed-shattering CGs from wheat, barley or rice that correspond to IWG HG4, the only relevant gene that we found was the *Arabidopsis Seuss (SEU)* gene (Balanza et al. 2016), which aligned to the SESH and SENA QTLs on LG10 and the SESH QTL on LG12. Many genes controlling photoperiod and flowering date have been identified in temperate cereals including wheat and barley (Cockram et al. 2007). The *vernalization 2 (VRN2)* gene is a major determinant of flowering time in wheat (Yan et al. 2004) and barley (Dubcovsky et al. 2005; Karsai et al. 2005; Laurie et al. 1995). The *VRN2* gene is located on barley chromosome 4HL and a region of wheat chromosome 4AL that was translocated to 5AL (Devos et al. 1995; Dubcovsky et al. 1998). The *VRN2* gene was located on IWG CHR10 (HG4) and CHR15 (HG5), which was not unexpected because LG12 (HG4) and LG15 (HG5) of IWG display a reciprocal translocation (Kantarski et al. 2017) similar to the 4AL/5AL translocation of *Triticum monococcum*, *T. aestivum* and other Triticeae species (Devos et al. 1995; Dubcovsky et al. 1998; Larson et al. 2012). A third *VRN2* gene was found on an unmapped scaffold\_264 that probably belongs to CHR11, which is colinear with barley chromosome 4H (Kantarski et al. 2017). However, none of the ZAMA QTLs aligned to the CHR10-, CHR15-, or predicted CHR11-*VRN2* loci. Two *Phytochrome* genes designated *PHYA* and *PHYB* (Mathews and Sharrock 1996) map to CHR10, CHR11, and CHR12 with alignments to ZAMA QTLs on LG10 and LG11. The LG10 ZAMA QTL could be a pleiotropic effect of the LG10 STLE QTL, but there was no alternative explanation for the alignment of the LG11 ZAMA QTL to the CHR10 *PHYB* gene.

Homoeologous group 5 was exceptional in that it displayed the greatest number of QTLs, 25 in total, and that it displayed significant effects for SEFL and SESP on all three HG5 linkage groups including LG13, LG14, and LG15 (Table 5). The *Q* gene on wheat chromosome 5AL confers the free-threshing trait of domesticated wheat and has pleiotropic effects on glume shape, glume tenacity, rachis fragility, spike length, plant height and heading date (Faris and Gill 2002; Simons et al. 2006). Orthologs of *Q* were found on the long arm of CHR13, CHR14, and CHR15 with alignment to INLE, SESH, and SENA QTLs on LG14 and possibly another SENA QTL on LG15 (Figure 1). Other potentially important CGs located near *Q* include the *Phytochrome C (PHYC)* and *Vernalization 1* (Yan et al. 2003) genes, which aligned to a ZAMA QTL on LG14 (Figure 1). Orthologs of the *Constans 3 (CO3)* *CONSTANS*-like flowering gene (Griffiths et al. 2003) also align to ZAMA QTLs on LG14 and LG15. Homologs of the rice *Grain length chromosome-3 QTL (GL3)* gene (Zhang et al. 2012b) were present at two loci on CHR13, CHR14, and CHR15 with alignments to seed-size QTLs on all three corresponding linkage groups (Figure 1). Two closely-linked homologs of the rice *Thousand-grain weight 6 (TGW6)* gene (Ishimaru et al. 2013), located on CHR15, also aligned to seed-size QTLs on LG15. Homologs of the maize *Barren inflorescence 2 (BIF2)* gene

(McSteen et al. 2007) and the *Dense and erect panicle 1 (DEP1)* gene of rice and barley (Huang et al. 2009; Wendt et al. 2016) show alignments to INLE and SPIN QTLs on LG13 and LG14.

Homoeologous group 6 was tied with HG2 for having the highest SEAR LOD score and all three HG6 linkage groups (LG16, LG17, and LG18) had significant ZAMA effects (Table 5). Three *CONSTANS*-like flowering orthogenes designated *CO2*, *CO5* and *CO7* (Griffiths et al. 2003) were each present on CHR16, CHR17, and CHR18 except that *CO5* was not present on CHR16. At least one of two of these *CONSTANS* genes aligned to each of the ZAMA QTLs on LG16 and LG17, but none aligned to the ZAMA QTL on the distal long arm of LG18. Orthologs of the *Growth-regulating factor 4 (GRF4)* gene (Sun et al. 2016), *Grain weight chromosome-2 QTL (GW2)* gene (Song et al. 2007; Su et al. 2011), and a *GDSL lipase (GLIP)* gene aligned to seed-size QTLs on LG16, LG17, and LG18. The rice *GRF4* gene encodes a growth and cytokinin-biosynthesis regulator that increases grain size and inflorescence length and decreases seed shattering (Sun et al. 2016). The *GLIP* locus on LG17 contained one of the most significant SELE QTL markers in the M26 x M35 family, TP678810, which was also associated with SEMA, SEAR, and SELE in the UMN AM population (Zhang et al. 2017). Some *GLIP* genes (Jiang et al. 2012) have roles in seed size and metabolism (Clauß et al. 2011; Huang et al. 2015; Lai et al. 2017; Ma et al. 2018; Tiwari et al. 2016).

Homoeologous group 7 was remarkable in that it was associated with a total of 21 seed-size QTLs, only one less than HG2, and the highest number of M26 x M35 seed-size QTLs (Table 5). Compared to the other HGs, more seed-size CGs and CG-QTL alignments were also found in HG7. Six relevant seed-size CGs including the rice chromosome-8 *Fertilization independent seed (FEI)* gene (Kapazoglou et al. 2010; Nallamilli et al. 2013), rice *Grain width chromosome-8 QTL (GW8)* gene (Wang et al. 2012), *Glucan, water-dikinase (GWD)* gene (Ral et al. 2012; Shu and Rasmussen 2014), *Sucrose synthase 1 (SUS1)* gene (Hou et al. 2014; Jiang et al. 2011), *Thousand-grain weight 6 (TGW6)* gene (Ishimaru et al. 2013), and wheat *Thousand-grain weight chromosome-7A (TGW7A)* gene (Hu et al. 2016) all aligned to seed-size QTLs in HG7 (Table 8). The maize *teosinte glume architecture 1 (tga1)* gene (Wang et al. 2005) and barley *nud* gene (Taketa et al. 2008), responsible for naked grains, were both found on LG19, LG20, and LG21 of HG7 but no significant IWG SENA QTLs were detected in HG7.

### **The S and Z self-incompatibility (SI) genes**

The S and Z SI genes have been shown to affect seed size and fertility traits in perennial ryegrass (*Lolium perenne*) if there is insufficient genetic variability in populations such as a full-sib mapping family (Studer et al. 2008), which raises concerns about possible effects of these genes in the full-sib IWG QTL mapping populations. The S and Z SI genes could also be a yield-limiting factor in the broader IWG grain breeding populations considering that they purportedly incurred population bottlenecks as few as 14 individuals (Wagoner 1990; Zhang et al. 2016). The S and Z genes are located on chromosomes that correspond to homoeologous groups 1 and 2, respectively, in both *Lolium* and *Secale* (Hackauf and Wehling 2005; Manzanares et al. 2016; Shinozuka et al. 2010; Thorogood et al. 2017). Seed fertility QTLs (SEFL and SESP) were detected on 10 linkage groups (Table 5) including LG01 (HG1) and LG06 (HG2). In fact, LG06 displayed the highest LOD scores for several seed-size (SEMA, SEAR, and SELE), seed-fertility (SEFL and SESP), and seed yield (SYIN) traits. Identification of S and Z genes has been difficult

(Thorogood et al. 2017), but it is thought that a *domain of unknown function* (*DUF247*) gene is the pollen component of the *S* locus on HG1 (Manzanares et al. 2016). Another paralogous *DUF247* gene (Shinozuka et al. 2010) and closely linked *ubiquitin-specific protease* (*USP*) gene (Hackauf and Wehling 2005) are considered the best candidates for the *Z* locus (Thorogood et al. 2017). Putative orthologs of the *S* (*DUF247*) gene were found on IWG CHR01, CHR02, and CHR03 of HG1 with alignments to seed-size QTLs (Figure 1). Moreover, two markers with significant seed-mass effects in the AM population (Zhang et al. 2017) were located about 5 MB and 68 KB from the LG01 and LG02 *S* loci, respectfully (Figure 1). However, none of the putative IWG *S* genes aligned to SESP or SEFL fertility QTLs (Figure 1). Tightly-linked *DUF247* and *USP* genes on IWG CHR01 (HG1) and CHR06 (HG2) were homologous to tightly-linked *DUF247* and *USP* genes corresponding to the *Z* ortholoci on HG2 of *Secale* (Hackauf and Wehling 2005) and *Lolium* (Shinozuka et al. 2010). The putative *Z* locus on IWG CHR06 shows possible alignment near seed-size, fertility, and seed-yield QTLs on LG06 and it is presumably orthologous to the *Z* locus on HG2 of *Secale* (Hackauf and Wehling 2005) and *Lolium* (Shinozuka et al. 2010). Linkage group 6 (LG06) was the only LG that had significant QTLs for all four seed-size traits in both M26 x M35 and C3-2331 x C3-2595 families and LG06 also had the highest LOD scores for SEFL and SESP (Table 5). However, it was also interesting that CHR06 was one of only three chromosomes that did not have any significant seed-size AM effects (Zhang et al. 2017) indicating that diversity of the putative IWG CHR07 *Z* locus was not limiting seed size in the AM population. Tightly linked *DUF247* and *USP* genes on IWG CHR01 aligned to SESP and SYIN QTLs on LG01 (Figure 1), but it is not absolutely clear if this CHR01 (HG1) IWG locus is orthologous to the HG2 *Z* locus of *Secale* and *Lolium* (Hackauf and Wehling 2005; Shinozuka et al. 2010). Further research is needed to determine if the *S* or *Z* SI genes are affecting grain size and grain yields in the IWG grain breeding populations and future experimental research should be designed to consider this question.

### Prospects for gene identification

Recent advancements in DNA sequencing and genomics have enabled scientists to pinpoint genes and chromosome regions that distinguish crops from their wild progenitors, with promising applications to a multitude of additional crop species (Tang et al. 2010). Development of high-density GBS linkage maps (Kantarski et al. 2017) and a draft genome sequence will facilitate identification of CGs associated with IWG domestication QTLs, but it is also recognized that most of these QTLs span large regions of the genome containing many other genes (Figure 1). Additional sequencing of M26, M36, C4-3471, and other IWG reference plants will facilitate identification of mutations in these CGs, some of which may have large or dramatic effects (Tang et al. 2010). Moreover, relatively low levels of linkage disequilibrium and high-levels of outcrossing coupled with recent advances in DNA sequencing and genotyping will facilitate genetic testing of CG variants by association mapping in IWG (Zhang et al. 2017). This approach was used to identify or test CGs of wheat (Hou et al. 2014; Hu et al. 2016; Jiang et al. 2011; Ma et al. 2016; Su et al. 2011; Zhang et al. 2012a), barley (Shu and Rasmussen 2014), maize (Gallavotti et al. 2004), and rice (Li et al. 2011; Wang et al. 2015; Weng et al. 2008; Zhang et al. 2012a) examined in this study (Table 8). However, it is likely that many of the IWG QTLs and AM effects are caused by other genes not examined in this study or perhaps not recognized in any other crop species. Positional cloning was used to identify many CGs of the wheat (Faris et al. 2003; Ishimaru et al. 2013; Yan et al. 2003; Yan et al. 2004), barley (Komatsuda et al. 2007; Vu

et al. 2010), rice (Ashikari et al. 2005; Huang et al. 2009; Li et al. 2011; Song et al. 2007; Sun et al. 2016; Wang et al. 2008; Wang et al. 2012; Wang et al. 2015; Weng et al. 2008; Wu et al. 2016; Zhang et al. 2012b), and *Lolium* (Manzanares et al. 2016; Shinozuka et al. 2010) examined in this study (Table 8). Positional cloning has been particularly successful in rice in part because of its relatively small genome size, but recent advances in DNA sequencing will accelerate progress in large-genome species including wheat, barley, and IWG.

Comparisons of QTL results from the M26 x M35 and C3-2331 x C3-2595 families identified possible targets for positional cloning. The M26 x M35 family was specifically developed to examine the unique combination seed-retention and naked-seed traits found in C3\_3471, by crossing C3\_3471 with a more primitive plant, C3\_3941, and testing for segregation of QTLs from the F1 hybrid parent, M26. Results of this experiment indicate that this unique combination of SESH and SENA traits, observed in C3\_3471 (parent of M26), involved a combination of at least two QTLs located on LG10 and LG11, which had synergistic M26- $\alpha$  effects on both traits (Table 6). The LG11 SESH QTL had the highest LOD score for any trait (Table 5) and it may be a reasonable target for positional cloning especially if this QTL could be isolated in an otherwise isogenic or neutral background. Other possible targets for positional cloning are the SENA QTL on LG02 and seed-size and seed-yield QTLs on LG06, but there are also some potentially good CGs for these QTLs that should be considered as discussed above. The LG02 SENA QTL on LG02 had relatively large effects as percentage of the mean (Table 6), meaning that it may have high breeding value. However, the LOD score of the LG02 SENA QTL was not so exceptional (Table 5), meaning that it may be difficult to obtain the mapping precision required for positional cloning unless measurements can be improved, or the experimental complexity can be reduced. Relatively strong seed-size and seed-yield QTLs were associated with LG06 in both M26 x M35 and C3-2331 x C3-2595 families (Figure 1, Table 5) and the LG06 SELE QTL was the only QTL that was significant across all five environments (Table 4), but additional research is needed to determine if these LG06 QTLs were caused by the *Z SI* gene or some other candidate gene such as *GIF*, *SDW3*, *SUS2*, or *GL7* (Figure 1) before positional cloning should be considered.

### **Implications for IWG domestication and improvement**

Transgressive genetic variation, where progeny (Table 2) exceed parents (Table 1), and 111 significant QTLs were observed for 17 traits related to seed size, reproductive fertility, inflorescence capacity, stem and inflorescence compactness, seed retention in the field, percentage of naked seeds after threshing, and maturity in the M26 x M35 family. Theoretically, fixation of the optimum QTL genotypes could improve trait means in the M26 x M35 family (Table 2) by an average of 12% to 118% across all five environments (Table 6) but this would require a uniform hybrid with one copy of the best M26 QTL allele and one copy of the best M35 allele for each QTL. It may be possible to make greater improvements by fixing the best M26 allele or the best M35 QTL allele, as homozygous genotypes, but the breeding values of these genotypes are unknown because all progeny contained one M26 allele (*a* or *b*) and one M35 allele (*c* or *d*). Thus, marker-assisted selection for most of these traits may be challenging considering the number and complexity of QTL effects and the fact that the M26 x M35 family is a small sample of the genetic variation present in genetically heterogeneous IWG grain breeding populations (Zhang et al. 2016). However, comparisons of QTLs and traits from the M26 x M35 family to those of fully domesticated grain crops, such as wheat, will elucidate pathways for successful IWG domestication.

Grain weights of wheat normally range from about 27 to 60 mg per grain among wheat landraces and cultivars (Abbo et al. 2014; Gegas et al. 2010; Ma et al. 2016) with up to 65 seeds per spike (Wang et al. 2010). Seed yield potential of wheat normally ranges from about 1,500 to 2,400 mg per spike, but up 5,000 mg per spike has been reported for Tibetan Triple-Spikelet wheats that have up to 121 seeds per spike (Yang et al. 2005). With up to 37 spikelets per inflorescence and four seeds per spikelet observed on some IWG genets, the total number of seeds possible on each spike (148) exceeds that of wheat or barley. With nearly 9 mg per seed on some M26 x M35 IWG genets (Table 2) and up to 12 or 14 mg per seed on others IWG plants (Cattani and Asselin 2018; Zhang et al. 2017), the maximum theoretical seed yield per spike would be nearly 2,100 mg. The best M26 x M35 genets produced no more than 525 mg of seed per spike, but up to and 932 mg of seed per spike has been reported for some IWG plants (Cattani 2017). With an average of about seven (Table 2) to nine (Cattani and Asselin 2018) florets per spikelet, up to 37 spikelets per head (Table 2), and grain weights of 12 to 14 mg the maximum theoretical seed yield per spike for IWG would be from 3,100 to 4,600 mg if every floret produced seed. However, we never observed more than 49% floret fertility in the M26 x M35 family (Table 2). Fertility is considered one of the limiting factors in grass seed production (Armstead et al. 2008) and we observed a relatively large number of QTLs with relatively large QTL effects for SEFL and SESP traits in the M26 x M35 family (Tables 5 and 6). These findings indicate that selection for greater fertility may be a promising avenue to improve IWG seed yields, with potential to match the seed-yield per spike of wheat, but domesticated crops often have fewer and larger fruits or grains compared to their progenitors (Doebley et al. 2006).

Seed retention, free-threshing (naked) seed, and increased seed size are all key domestication or improvement traits of most grain crops (Abbo et al. 2014; Doebley et al. 2006; Gegas et al. 2010; Kovach et al. 2007; Lenser and Theißen 2013; Liu et al. 2016; Meyer and Purugganan 2013) including IWG. Compared to other traits evaluated in this experiment, IWG seed shattering (SESH) was controlled by a relatively small number of QTLs with relatively large QTL effects (Table 6), including the highest LOD score in this study (Table 5), which is similar to observations in other grasses and grain crops (Doebley et al. 2006; Doust et al. 2014; Kovach et al. 2007; Larson and Kellogg 2009; Pourkheirandish et al. 2015; Simons et al. 2006). Conversely, genetic factors controlling seed-threshing (SENA), seed-size (SEMA, SEAR, SELE, and SEWI), and other traits such as maturity (ZAMA) were substantially more complex. Compared to other traits, SENA and ZAMA were controlled by relatively large numbers of QTLs (Tables 5 and 6) that explained more phenotypic variation in the M26 x M35 family (Table 7). Although we only detected four SEMA QTLs, there were a total of at least 12 loci affecting at least one of the four seed-size traits (SEMA, SEAR, SELE, and SEWI). Interestingly, the relative magnitudes of Avg-total QTL effects (Table 6) and PVE (Table 7) for SENA and SEMA were similar to the expected selection gains of 181% and 60% over five cycles of selection for these two traits, respectively (DeHaan et al. 2018). The average seed weights of IWG forage varieties, ranging from about 2 to 8 mg per seed (Berdahl and Frank 1998; Schulz-Schaeffer and Haller 1987), overlaps with the associated with IWG plants selected for grain production, which range from 3 to 14 mg per seed as reported in this and other studies (Cattani and Asselin 2018; Zhang et al. 2017). Similar overlap in seed sizes exists among wild and domestic forms of barley, wheat, and rice (Abbo et al. 2014; Fuller 2007; Gegas et al. 2010), but kernel weight has reportedly increased 10-fold during the domestication of maize (Liu et al. 2016). Evidence

from this and other studies demonstrate that IWG seed size can be improved with some rapid initial gains (Cattani and Asselin 2018; DeHaan et al. 2018; Zhang et al. 2017), but additional improvements in seed size may be slow and incremental or require new discoveries and breakthroughs. Conversely, relatively strong QTL effects for SENA and SESH, discussed above, suggest that free-threshing (naked) seed and strong seed retention are attainable domestication traits (Abbo et al. 2014; Doebley et al. 2006) that will clearly distinguish wild and domestic forms of IWG.

## Conclusions

A total of 111 QTLs were detected for 17 variable traits in the M26 x M35 family including several large-effect QTLs responsible for critical IWG domestication and improvement traits related to fertility, inflorescence architecture, plant height, seed retention, seed size and seed-threshing. The magnitude of M26 x M35 QTL effects, heritabilities and range of phenotypic variation observed in this and other studies (Cattani and Asselin 2018; Zhang et al. 2017) demonstrate potential to fix critical domestication traits, including seed retention and free-threshing (naked) seed, and improve other important grain production traits in IWG. With up to four possible alleles for each M26 x M35 QTL and even greater complexity in genetically heterogeneous breeding populations, the prospects for marker-assisted selection are not certain. However, identification of genes and loci directly associated with critical IWG domestication and improvement traits will enable better management and utilization of IWG germplasm. A total 42 domestication orthogenes, including the wheat free-threshing *Q* (Simons et al. 2006) and *reduced-height* green revolution (Peng et al. 1999) genes, aligned to one or more relevant QTLs in this experiment. Closely-linked *DUF247* and *USP* genes on IWG CHR06 corresponding to the *Z* self-incompatibility locus of *Secale* and *Lolium* (Hackauf and Wehling 2005; Shinozuka et al. 2010) showed possible alignment to seed-size, fertility, and seed-yield QTLs on IWG LG06 suggesting that diversity of SI genes may be a limiting factor for seed production in the full-sib IWG families. A large-effect QTL, with a LOD score of nearly 15 for seed shattering, had synergistic effects resulting in greater seed retention and more free-threshing (naked) seed in the M26 x M35 family. Although no candidate genes were associated with this large-effect QTL, recent advancements in genome sequencing and genotyping provide useful approaches to pinpoint genes or loci responsible for domestication of other crops (Tang et al. 2010) such as IWG. Herein, two DNA markers with relatively strong effects on IWG seed size across independent QTL and association mapping experiments (Zhang et al. 2017) were located directly within IWG *GDSL* *lipase* (Clauß et al. 2011; Huang et al. 2015; Jiang et al. 2012; Lai et al. 2017; Ma et al. 2018; Tiwari et al. 2016) and *Xylogalacturonan (XGA) xylosyltransferase* (Jensen et al. 2008; Zhang et al. 2015) orthogenes, which may be plausible candidate genes for these IWG seed-size QTLs. Results of this study demonstrated the increasing power of high-density genotyping (Kantarski et al. 2017), genome sequencing, and QTL mapping to elucidate pathways for the domestication and improvement of a new and genetically-complex perennial grain crop.

## Author Contribution Statement

SL and LD conceived M26 x M35 QTL experiment and conducted field evaluations. SL performed QTL analyses and wrote manuscript. JP and TK genotyped the M26 x M35 plants. XZ and JA conducted C3-2331 x C3-2595 QTL

and AM experiments and provided unpublished data from these experiments. TK, SL, XZ, LD, JA, and JP developed linkage maps used for all QTL and AM analyses. JP, LD, KD, and SL conceived the IWG genome sequencing project as described in JGI Proposal Id. 1997. JS, JG, JJ, and SS assembled and annotated the IWG draft genome sequence. JC and MR assisted with data management and data analysis. KJ provided guidance and support for field and trait evaluations in Utah. All authors read and reviewed the manuscript.

## Acknowledgments

This research was supported by The Malone Family Land Preservation Foundation. The work conducted by the US Department of Energy Joint Genome Institute is supported by the Office of Science of the US Department of Energy under Contract No. DE-AC02-05CH11231. KD is supported by USDA-NIFA Post Doctoral Fellowships grant no. 2017-67012-26129/project accession no. 1011622 “Exploring the Genomic Landscape of Perenniality within the Triticeae”. The authors wish to thank Martin Mascher and LiangLiang Gao for advice and contributions to the continuing development of the *Thinopyrum intermedium* genome assembly.

## Conflict of Interest

On behalf of all authors, the corresponding author states that there is no conflict of interest.

## References

- (!!! INVALID CITATION !!! (Manzanares et al. 2016; Thorogood et al. 2017)).
- Abbo S, Pinhasi van-Oss R, Gopher A, Saranga Y, Ofner I, Peleg Z (2014) Plant domestication versus crop evolution: a conceptual framework for cereals and grain legumes. Trends Plant Sci 19:351-360. doi:<https://doi.org/10.1016/j.tplants.2013.12.002>
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. J Mol Biol 215:403-410. doi:10.1016/S0022-2836(05)80360-2
- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res 25:3389-3402. doi:10.1093/nar/25.17.3389
- Armstead IP, Turner LB, Marshall AH, Humphreys MO, King IP, Thorogood D (2008) Identifying genetic components controlling fertility in the outcrossing grass species perennial ryegrass (*Lolium perenne*) by quantitative trait loci analysis and comparative genetics. New Phytol 178:559-571. doi:10.1111/j.1469-8137.2008.02413.x
- Ashikari M et al. (2005) Cytokinin oxidase regulates rice grain production. Science 309:741-745. doi:10.1126/science.1113373
- Balanza V, Roig-Villanova I, Di Marzo M, Masiero S, Colombo L (2016) Seed abscission and fruit dehiscence required for seed dispersal rely on similar genetic networks. Development 143:3372-3381. doi:10.1242/dev.135202

787 Bell LW, Harrison MT, Kirkegaard JA (2015) Dual-purpose cropping – capitalising on potential grain crop grazing  
 788 to enhance mixed-farming profitability. *Crop and Pasture Science* 66:i-iv. doi:10.1071/CPv66n4\_FO  
 789 Berdahl JD, Frank AB (1998) Seed maturity in four cool-season forage grasses. *Agron J* 90:483-488  
 790 Cattani D, Asselin S (2018) Has Selection for Grain Yield Altered Intermediate Wheatgrass? *Sustainability* 10:688  
 791 Cattani DJ (2017) Selection of a perennial grain for seed productivity across years: Intermediate wheatgrass as a test  
 792 species. *Can J Plant Sci* 97:516-524. doi:10.1139/cjps-2016-0280  
 793 Cattani DJ, Asselin SR (2017) Extending the growing season: forage seed production and perennial grains. *Can J*  
 794 *Plant Sci* 98:235-246. doi:10.1139/cjps-2017-0212  
 795 Ceoloni C, Kuzmanovic L, Forte P, Virili ME, Bitti A (2015) Wheat-perennial Triticeae introgressions: Major  
 796 achievements and prospects. In: *Alien Introgression in Wheat: Cytogenetics, Molecular Biology, and*  
 797 *Genomics*. pp 273-313. doi:10.1007/978-3-319-23494-6-11  
 798 Chen G, Li H, Wei Y, Zheng Y-L, Zhou M, Liu C (2016) Pleiotropic effects of the semi-dwarfing gene *uzu* in  
 799 barley. *Euphytica* 209:749-755. doi:10.1007/s10681-016-1668-4  
 800 Chono M et al. (2003) A semidwarf phenotype of barley *uzu* results from a nucleotide substitution in the gene  
 801 encoding a putative brassinosteroid receptor. *Plant Physiol* 133:1209-1219. doi:10.1104/pp.103.026195  
 802 Clauß K et al. (2011) Overexpression of sinapine esterase *BnSCE3* in oilseed rape seeds triggers global changes in  
 803 seed metabolism. *Plant Physiol* 155:1127-1145. doi:10.1104/pp.110.169821  
 804 Cockram J, Jones H, Leigh FJ, O'Sullivan D, Powell W, Laurie DA, Greenland AJ (2007) Control of flowering time  
 805 in temperate cereals: genes, domestication, and sustainable productivity. *J Exp Bot* 58:1231-1244  
 806 Cox TS, Glover JD, van Tassel DL, Cox CM, DeHaan LR (2006) Prospects for developing perennial grain crops.  
 807 *Bioscience* 56:649-659  
 808 Cox TS, Van Tassel DL, Cox C, DeHaan L (2010) Progress in breeding perennial grains. *Crop and Pasture Science*  
 809 61:513-521. doi:10.1071/CP09201  
 810 Culman SW, Snapp SS, Ollenburger M, Basso B, DeHaan LR (2013) Soil and water quality rapidly responds to the  
 811 perennial grain Kernza wheatgrass. *Agron J* 105:735-744. doi:10.2134/agronj2012.0273  
 812 Curwen-McAdams C, Jones SS (2017) Breeding perennial grain crops based on wheat. *Crop Sci* 57:1172-1188.  
 813 doi:10.2135/cropsci2016.10.0869  
 814 DeHaan L, Christians M, Crain J, Poland J (2018) Development and evolution of an intermediate wheatgrass  
 815 domestication program. *Sustainability* 10:1499  
 816 DeHaan LR, Ismail BP (2017) Perennial cereals provide ecosystem benefits. *Cereal Foods World* 62:278-281.  
 817 doi:10.1094/CFW-62-6-0278  
 818 DeHaan LR et al. (2016) A pipeline strategy for grain crop domestication. *Crop Sci* 56:917-930.  
 819 doi:10.2135/cropsci2015.06.0356  
 820 Devos KM, Dubcovsky J, Dvorak J, Chinoy CN, Gale MD (1995) Structural evolution of wheat chromosomes 4A,  
 821 5A, and 7B and its impact on recombination. *Theor Appl Genet* 91:282-288. doi:10.1007/BF00220890  
 822 Doebley J, Stec A, Hubbard L (1997) The evolution of apical dominance in maize. *Nature* 386:485-488.  
 823 doi:10.1038/386485a0



824 Doebley JF, Gaut BS, Smith BD (2006) The Molecular Genetics of Crop Domestication. *Cell* 127:1309-1321.  
825 doi:<https://doi.org/10.1016/j.cell.2006.12.006>

826 Doust AN, Mauro-Herrera M, Francis AD, Shand LC (2014) Morphological Diversity and Genetic Regulation of  
827 Inflorescence Abscission Zones in Grasses. *Am J Bot* 101:1759-1769

828 Dubcovsky J, Chen CL, Yan LL (2005) Molecular characterization of the allelic variation at the *VRN-H2*  
829 vernalization locus in barley. *Mol Breed* 15:395-407. doi:10.1007/s11032-005-0084-6

830 Dubcovsky J, Lijavetzky D, Appendino L, Tranquilli G (1998) Comparative RFLP mapping of *Triticum*  
831 *monococcum* genes controlling vernalization requirement. *Theor Appl Genet* 97:968-975.  
832 doi:10.1007/s001220050978

833 Faris JD, Fellers JP, Brooks SA, Gill BS (2003) A bacterial artificial chromosome contig spanning the major  
834 domestication locus Q in wheat and identification of a candidate gene. *Genetics* 164:311-321

835 Faris JD, Gill BS (2002) Genomic targeting and high-resolution mapping of the domestication gene *Q* in wheat.  
836 *Genome* 45:706-718

837 Fuller DQ (2007) Contrasting Patterns in Crop Domestication and Domestication Rates: Recent Archaeobotanical  
838 Insights from the Old World. *Ann Bot* 100:903-924. doi:10.1093/aob/mcm048

839 Gallavotti A et al. (2004) The role of *barren stalk1* in the architecture of maize. *Nature* 432:630.  
840 doi:10.1038/nature03148

841 Galli M et al. (2015) Auxin signaling modules regulate maize inflorescence architecture. *Proc Natl Acad Sci USA*  
842 112:13372-13377. doi:10.1073/pnas.1516473112

843 Gegas VC et al. (2010) A Genetic Framework for Grain Size and Shape Variation in Wheat. *The Plant Cell* 22:1046-  
844 1056. doi:10.1105/tpc.110.074153

845 Griffiths S, Dunford RP, Coupland G, Laurie DA (2003) The evolution of *CONSTANS*-like gene families in barley,  
846 rice, and *Arabidopsis*. *Plant Physiol* 131:1855-1867. doi:10.1104/pp.102.016188

847 Hackauf B, Wehling P (2005) Approaching the self-incompatibility locus Z in rye (*Secale cereale* L.) via  
848 comparative genetics. *Theor Appl Genet* 110:832-845. doi:10.1007/s00122-004-1869-4

849 Harmony KR (2015) Cool-season grass biomass in the southern mixed-grass prairie region of the USA. *Bioenergy*  
850 *Research* 8:203-210. doi:10.1007/s12155-014-9514-9

851 Hayes R et al. (2018) The performance of early-generation perennial winter cereals at 21 sites across four  
852 continents. *Sustainability* 10:1124

853 Holland JB, Nyquist WE, Cervantes-Martínez CT (2010) Estimating and Interpreting Heritability for Plant  
854 Breeding: An Update. *Plant Breeding Reviews*. doi:10.1002/9780470650202.ch2

855 Hou J, Jiang Q, Hao C, Wang Y, Zhang H, Zhang X (2014) Global selection on sucrose synthase haplotypes during  
856 a century of wheat breeding. *Plant Physiol* 164:1918-1929. doi:10.1104/pp.113.232454

857 Houston K et al. (2013) Variation in the interaction between alleles of *HvAPETALA2* and microRNA172 determines  
858 the density of grains on the barley inflorescence. *Proc Natl Acad Sci USA* 110:16675-16680.  
859 doi:10.1073/pnas.1311681110

860 Hu M-J et al. (2016) Cloning and characterization of *TaTGW-7A* gene associated with grain weight in wheat via  
 861 SLAF-seq-BSA. *Front Plant Sci* 7:1902. doi:10.3389/fpls.2016.01902  
 862 Huang L-M, Lai C-P, Chen L-FO, Chan M-T, Shaw J-F (2015) *Arabidopsis SFAR4* is a novel GDSL-type esterase  
 863 involved in fatty acid degradation and glucose tolerance. *Bot Stud* 56:33. doi:10.1186/s40529-015-0114-6  
 864 Huang X et al. (2009) Natural variation at the *DEP1* locus enhances grain yield in rice. *Nat Genet* 41:494-497.  
 865 doi:10.1038/ng.352  
 866 Ishimaru K et al. (2013) Loss of function of the IAA-glucose hydrolase gene *TGW6* enhances rice grain weight and  
 867 increases yield. *Nat Genet* 45:707. doi:10.1038/ng.2612  
 868 Jensen JK et al. (2008) Identification of a Xylogalacturonan Xylosyltransferase Involved in Pectin Biosynthesis in  
 869 *Arabidopsis*. *The Plant Cell* 20:1289-1302. doi:10.1105/tpc.107.050906  
 870 Jensen KB, Yan X, Larson SR, Wang RRC, Robins JG, McIntosh RA (2016) Agronomic and genetic diversity in  
 871 intermediate wheatgrass (*Thinopyrum intermedium*). *Plant Breeding* 135:751-758. doi:10.1111/pbr.12420  
 872 Jensen KB, Zhang YF, Dewey DR (1990) Mode of Pollination of Perennial Species of the Triticeae in Relation to  
 873 Genomically Defined Genera. *Can J Plant Sci* 70:215-225  
 874 Jia QJ, Zhang JJ, Westcott S, Zhang XQ, Bellgard M, Lance R, Li CD (2009) GA-20 oxidase as a candidate for the  
 875 semidwarf gene *sdw1/denso* in barley. *Functional & Integrative Genomics* 9:255-262  
 876 Jiang Q, Hou J, Hao C, Wang L, Ge H, Dong Y, Zhang X (2011) The wheat (*T. aestivum*) sucrose synthase 2 gene  
 877 (*TaSus2*) active in endosperm development is associated with yield traits. *Functional & Integrative*  
 878 *Genomics* 11:49-61. doi:10.1007/s10142-010-0188-x  
 879 Jiang Y, Chen R, Dong J, Xu Z, Gao X (2012) Analysis of GDSL lipase (GLIP) family genes in rice (*Oryza sativa*).  
 880 *Plant OMICS* 5:351-358  
 881 Jungers JM, DeHaan LR, Betts KJ, Sheaffer CC, Wyse DL (2017) Intermediate wheatgrass grain and forage yield  
 882 responses to nitrogen fertilization. *Agron J* 109:462-472. doi:10.2134/agronj2016.07.0438  
 883 Kantarski T, Larson S, Zhang X, DeHaan L, Borevitz J, Anderson J, Poland J (2017) Development of the first  
 884 consensus genetic map of intermediate wheatgrass (*Thinopyrum intermedium*) using genotyping-by-  
 885 sequencing. *Theor Appl Genet* 130:137-150. doi:10.1007/s00122-016-2799-7  
 886 Kapazoglou A et al. (2010) Epigenetic chromatin modifiers in barley: IV. The study of barley *Polycomb* group  
 887 (*PcG*) genes during seed development and in response to external ABA. *BMC Plant Biol* 10:73.  
 888 doi:10.1186/1471-2229-10-73  
 889 Karsai I et al. (2005) The *Vrn-H2* locus is a major determinant of flowering time in a facultative x winter growth  
 890 habit barley (*Hordeum vulgare* L.) mapping population. *Theor Appl Genet* 110:1458-1466.  
 891 doi:10.1007/s00122-005-1979-7  
 892 Kenneth PV, Kevin JJ (2001) Adaptation of perennial Triticeae to the eastern Central Great Plains. *Journal of Range*  
 893 *Management* 54:674-679. doi:10.2307/4003670  
 894 Klaas M et al. (2011) Progress towards elucidating the mechanisms of self-incompatibility in the grasses: further  
 895 insights from studies in *Lolium*. *Ann Bot* 108:677-685. doi:10.1093/aob/mcr186

896 Knowles RP (1977) Recurrent mass selection for improved seed yields in intermediate wheatgrass. *Crop Sci* 17:51-  
897 54. doi:10.2135/cropsci1977.0011183X001700010015x

898 Komatsuda T et al. (2007) Six-rowed barley originated from a mutation in a homeodomain-leucine zipper I-class  
899 homeobox gene. *Proc Natl Acad Sci USA* 104:1424-1429

900 Konishi S, Izawa T, Lin SY, Ebana K, Fukuta Y, Sasaki T, Yano M (2006) An SNP caused loss of seed shattering  
901 during rice domestication. *Science* 312:1392-1396. doi:10.1126/science.1126410

902 Koppolu R et al. (2013) *Six-rowed spike4 (Vrs4)* controls spikelet determinacy and row-type in barley. *Proc Natl*  
903 *Acad Sci USA* 110:13198-13203. doi:10.1073/pnas.1221950110

904 Kovach MJ, Sweeney MT, McCouch SR (2007) New insights into the history of rice domestication. *Trends Genet*  
905 23:578-587. doi:<https://doi.org/10.1016/j.tig.2007.08.012>

906 Krupinsky JM, Berdahl JD (2000) Selecting resistance to *Bipolaris sorokiniana* and *Fusarium graminearum* in  
907 intermediate wheatgrass. *Plant Dis* 84:1299-1302. doi:10.1094/PDIS.2000.84.12.1299

908 Kuczyńska A, Mikołajczak K, Ćwiek H (2014) Pleiotropic effects of the *sdw1* locus in barley populations  
909 representing different rounds of recombination. *Electron J Biotechnol* 17:217-223.  
910 doi:10.1016/j.ejbt.2014.07.005

911 Kumar S, You FM, Cloutier S (2012) Genome wide SNP discovery in flax through next generation sequencing of  
912 reduced representation libraries. *BMC Genomics* 13:684. doi:10.1186/1471-2164-13-684

913 La Rota M, Sorrells ME (2004) Comparative DNA sequence analysis of mapped wheat ESTs reveals the complexity  
914 of genome relationships between rice and wheat. *Funct Integr Genomics* 4:34-46. doi:10.1007/s10142-003-  
915 0098-2

916 Lai Y et al. (2017) Association mapping of grain weight, length and width in barley (*Hordeum vulgare*) breeding  
917 germplasm. *Int J Agric Biol* 19:1175-1186. doi:10.17957/IJAB/15.0406

918 Lang T et al. (2018) Precise identification of wheat – *Thinopyrum intermedium* translocation chromosomes carrying  
919 resistance to wheat stripe rust in line Z4 and its derived progenies. *Genome* 61:177-185. doi:10.1139/gen-  
920 2017-0229

921 Larson S et al. (2017) Development and testing of cool-season grass species, varieties and hybrids for biomass  
922 feedstock production in western North America. *Agronomy* 7:3

923 Larson SR, Kellogg EA (2009) Genetic dissection of seed production traits and identification of a major-effect seed  
924 retention QTL in hybrid *Leymus* (Triticeae) wildryes. *Crop Sci* 49:29-40. doi:10.2135/cropsci2008.05.0277

925 Larson SR et al. (2012) *Leymus* EST linkage maps identify 4NsL-5NsL reciprocal translocation, wheat-*Leymus*  
926 chromosome introgressions, and functionally important gene loci. *Theor Appl Genet* 124:189-206.  
927 doi:10.1007/s00122-011-1698-1

928 Laurie DA, Pratchett N, Snape JW, Bezant JH (1995) RFLP mapping of five major genes and eight quantitative trait  
929 loci controlling flowering time in a winter x spring barley (*Hordeum vulgare* L.) cross. *Genome* 38:575-  
930 585

931 Lee D, Owens VN, Boe A, Koo BC (2009) Biomass and seed yields of big bluestem, switchgrass, and intermediate  
 932 wheatgrass in response to manure and harvest timing at two topographic positions. *Glob Change Biol*  
 933 *Bioenergy* 1:171-179. doi:10.1111/j.1757-1707.2009.01008.x  
 934 Lenser T, Theißen G (2013) Molecular mechanisms involved in convergent crop domestication. *Trends Plant Sci*  
 935 18:704-714. doi:<https://doi.org/10.1016/j.tplants.2013.08.007>  
 936 Li B et al. (2013) Constitutive expression of cell wall invertase genes increases grain yield and starch content in  
 937 maize. *Plant Biotechnol J* 11:1080-1091. doi:10.1111/pbi.12102  
 938 Li C, Zhou A, Sang T (2006) Rice domestication by reducing shattering. *Science* 311:1936-1939  
 939 Li J et al. (2017) Introduction of *Thinopyrum intermedium* ssp. *trichophorum* chromosomes to wheat by trigenic  
 940 hybridization involving *Triticum*, *Secale* and *Thinopyrum* genera. *Planta* 245:1121-1135.  
 941 doi:10.1007/s00425-017-2669-9  
 942 Li W, Gill BS (2006) Multiple genetic pathways for seed shattering in the grasses. *Funct Integr Genomics* 6:300-  
 943 309. doi:10.1007/s10142-005-0015-y  
 944 Li Y et al. (2011) Natural variation in *GS5* plays an important role in regulating grain size and yield in rice. *Nat*  
 945 *Genet* 43:1266. doi:10.1038/ng.977  
 946 Liu H et al. (2017) Production and molecular cytogenetic characterization of a durum wheat-*Thinopyrum elongatum*  
 947 7E disomic addition line with resistance to *Fusarium* head blight. *Cytogenet Genome Res* 153:165-173  
 948 Liu Z, Garcia A, McMullen MD, Flint-Garcia SA (2016) Genetic analysis of kernel traits in maize-teosinte  
 949 introgression populations. *G3: Genes, Genomes, Genetics* 6:2523-2530. doi:10.1534/g3.116.030155  
 950 Ma L, Li T, Hao C, Wang Y, Chen X, Zhang X (2016) *TaGS5-3A*, a grain size gene selected during wheat  
 951 improvement for larger kernel and yield. *Plant Biotechnol J* 14:1269-1280. doi:10.1111/pbi.12492  
 952 Ma R, Yuan H, An J, Hao X, Li H (2018) A *Gossypium hirsutum* GDSL lipase/hydrolase gene (*GhGLIP*) appears to  
 953 be involved in promoting seed growth in *Arabidopsis*. *PLoS ONE* 13. doi:10.1371/journal.pone.0195556  
 954 Manzanares C et al. (2016) A gene encoding a DUF247 domain protein cosegregates with the *S* self-incompatibility  
 955 locus in perennial ryegrass. *Mol Biol Evol* 33:870-884. doi:10.1093/molbev/msv335  
 956 Marti A, Bock JE, Pagani MA, Ismail B, Seetharaman K (2016) Structural characterization of proteins in wheat  
 957 flour doughs enriched with intermediate wheatgrass (*Thinopyrum intermedium*) flour. *Food Chem* 194:994-  
 958 1002. doi:10.1016/j.foodchem.2015.08.082  
 959 Mathews S, Sharrock RA (1996) The phytochrome gene family in grasses (Poaceae): A phylogeny and evidence that  
 960 grasses have a subset of the loci found in dicot angiosperms. *Mol Biol Evol* 13:1141-1150.  
 961 doi:10.1093/oxfordjournals.molbev.a025677  
 962 McSteen P, Malcomber S, Skirpan A, Lunde C, Wu X, Kellogg E, Hake S (2007) *barren inflorescence2* Encodes a  
 963 co-ortholog of the PINOID serine/threonine kinase and is required for organogenesis during inflorescence  
 964 and vegetative development in maize. *Plant Physiol* 144:1000-1011. doi:10.1104/pp.107.098558  
 965 Meyer RS, Purugganan MD (2013) Evolution of crop species: genetics of domestication and diversification. *Nature*  
 966 *Reviews Genetics* 14:840. doi:10.1038/nrg3605  
 967 <https://www.nature.com/articles/nrg3605#supplementary-information>

968 Monono EM, Nyren PE, Berti MT, Pryor SW (2013) Variability in biomass yield, chemical composition, and  
 969 ethanol potential of individual and mixed herbaceous biomass species grown in North Dakota. *Ind Crops*  
 970 *Prod* 41:331-339. doi:10.1016/j.indcrop.2012.04.051  
 971 Nadolska-Orczyk A, Rajchel IK, Orczyk W, Gasparis S (2017) Major genes determining yield-related traits in wheat  
 972 and barley. *Theor Appl Genet* 130:1081-1098. doi:10.1007/s00122-017-2880-x  
 973 Nallamilli BRR, Zhang J, Mujahid H, Malone BM, Bridges SM, Peng Z (2013) Polycomb group gene *OsFIE2*  
 974 regulates rice (*Oryza sativa*) seed development and grain filling via a mechanism distinct from *Arabidopsis*.  
 975 *PLoS Genet* 9:e1003322. doi:10.1371/journal.pgen.1003322  
 976 Namikawa S, Kawakami Z (1934) On the occurrence of the haploid, triploid, and tetraploid plants in twin seedlings  
 977 of common wheat. *Proc Imp Acad Jap* 10:668-671  
 978 Ouellette LA, Reid RW, Blanchard SG, Brouwer CR (2018) LinkageMapView—rendering high-resolution linkage  
 979 and QTL maps. *Bioinformatics* 34:306-307. doi:10.1093/bioinformatics/btx576  
 980 Pearson CH, Larson SR, Keske CMH, Jensen KB (2015) Native grasses for biomass production at high elevations. .  
 981 In: Cruz VMV, Dierig DA (eds) *Handbook of plant breeding*. vol [9]. Springer, New York, pp 101-132  
 982 Peng JR et al. (1999) 'Green revolution' genes encode mutant gibberellin response modulators. *Nature* 400:256-261  
 983 Pourkheirandish M et al. (2015) Evolution of the grain dispersal system in barley. *Cell* 162:527-539.  
 984 doi:10.1016/j.cell.2015.07.002  
 985 R Core Team (2017) R: A Language and Environment for Statistical Computing. R Foundation for Statistical  
 986 Computing. Vienna, Austria. URL <https://www.R-project.org/>  
 987 Ral JP et al. (2012) Down-regulation of Glucan, Water-Dikinase activity in wheat endosperm increases vegetative  
 988 biomass and yield. *Plant Biotechnol J* 10:871-882. doi:10.1111/j.1467-7652.2012.00711.x  
 989 Revelle W (2018) psych: Procedures for psychological, psychometric, and personality research. Northwestern  
 990 University. Evanston, Illinois, USA. URL <https://CRAN.R-project.org/package=psych>  
 991 Robins JG (2010) Cool-season grasses produce more total biomass across the growing season than do warm-season  
 992 grasses when managed with an applied irrigation gradient. *Biomass & Bioenergy* 34:500-505.  
 993 doi:10.1016/j.biombioe.2009.12.015  
 994 Ross JG (1963) Registration of Oahe Intermediate Wheatgrass (Reg. No. 5). *Crop Sci* 3:373-373.  
 995 doi:10.2135/cropsci1963.0011183X000300040046x  
 996 Ryan MR, Crews TE, Culman SW, DeHaan LR, Hayes RC, Jungers JM, Bakker MG (2018) Managing for  
 997 multifunctionality in perennial grain crops. *Bioscience* 68:294-304. doi:10.1093/biosci/biy014  
 998 Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. *Nat Methods*  
 999 9:671-675  
 1000 Schulz-Schaeffer J, Haller SE (1987) Registration of Montana-2 Perennial XAgrotriticum Intermediodurum  
 1001 Khizhnyak. *Crop Sci* 27:822-823. doi:10.2135/cropsci1987.0011183X002700040058x  
 1002 Shinozuka H, Cogan NO, Smith KF, Spangenberg GC, Forster JW (2010) Fine-scale comparative genetic and  
 1003 physical mapping supports map-based cloning strategies for the self-incompatibility loci of perennial  
 1004 ryegrass (*Lolium perenne* L.). *Plant Mol Biol* 72:343-355. doi:10.1007/s11103-009-9574-y

1005 Shomura A, Izawa T, Ebana K, Ebitani T, Kanegae H, Konishi S, Yano M (2008) Deletion in a gene associated with  
 1006 grain size increased yields during rice domestication. *Nat Genet* 40:1023-1028. doi:10.1038/ng.169  
 1007 Shu X, Rasmussen S (2014) Quantification of amylose, amylopectin, and  $\beta$ -glucan in search for genes controlling  
 1008 the three major quality traits in barley by genome-wide association studies. *Front Plant Sci* 5:197.  
 1009 doi:10.3389/fpls.2014.00197  
 1010 Simons KJ, Fellers JP, Trick HN, Zhang ZC, Tai YS, Gill BS, Faris JD (2006) Molecular characterization of the  
 1011 major wheat domestication gene *Q*. *Genetics* 172:547-555. doi:10.1534/genetics.105.044727  
 1012 Song XJ, Huang W, Shi M, Zhu MZ, Lin HX (2007) A QTL for rice grain width and weight encodes a previously  
 1013 unknown RING-type E3 ubiquitin ligase. *Nat Genet* 39:623-630. doi:10.1038/ng2014  
 1014 Studer B, Jensen LB, Hentrup S, Brazauskas G, Kölliker R, Lübberstedt T (2008) Genetic characterisation of seed  
 1015 yield and fertility traits in perennial ryegrass (*Lolium perenne* L.). *Theor Appl Genet* 117:781-791.  
 1016 doi:10.1007/s00122-008-0819-y  
 1017 Su Z, Hao C, Wang L, Dong Y, Zhang X (2011) Identification and development of a functional marker of *TaGW2*  
 1018 associated with grain weight in bread wheat (*Triticum aestivum* L.). *Theor Appl Genet* 122:211-223.  
 1019 doi:10.1007/s00122-010-1437-z  
 1020 Sun PY et al. (2016) *OsGRF4* controls grain shape, panicle length and seed shattering in rice. *J Integr Plant Biol*  
 1021 58:836-847. doi:10.1111/jipb.12473  
 1022 Taketa S et al. (2008) Barley grain with adhering hulls is controlled by an ERF family transcription factor gene  
 1023 regulating a lipid biosynthesis pathway. *Proc Natl Acad Sci USA* 105:4062-4067.  
 1024 doi:10.1073/pnas.0711034105  
 1025 Tanabata T, Shibaya T, Hori K, Ebana K, Yano M (2012) SmartGrain: High-throughput phenotyping software for  
 1026 measuring seed shape through image analysis. *Plant Physiol* 160:1871-1880. doi:10.1104/pp.112.205120  
 1027 Tang H, Sezen U, Paterson AH (2010) Domestication and plant genomes. *Curr Opin Plant Biol* 13:160-166.  
 1028 doi:<https://doi.org/10.1016/j.pbi.2009.10.008>  
 1029 Thorogood D et al. (2017) A novel multivariate approach to phenotyping and association mapping of multi-locus  
 1030 gametophytic self-incompatibility reveals *S*, *Z*, and other loci in a perennial ryegrass (*Poaceae*) population.  
 1031 *Front Plant Sci* 8. doi:10.3389/fpls.2017.01331  
 1032 Tiwari GJ, Chiang MY, De Silva JR, Song BK, Lau YL, Rahman S (2016) Lipase genes expressed in rice bran:  
 1033 LOC\_Os11g43510 encodes a novel rice lipase. *J Cereal Sci* 71:43-52. doi:10.1016/j.jcs.2016.07.008  
 1034 Tulpan D, Leger S (2017) The Plant Orthology Browser: An Orthology and Gene-Order Visualizer for Plant  
 1035 Comparative Genomics. *The Plant Genome* 10. doi:10.3835/plantgenome2016.08.0078  
 1036 Uzma, Kubra G, Gul A, Mujeeb-Kazi A (2015) Use of alien diversity to combat some major biotic stresses in  
 1037 *Triticum aestivum* L. In: Hakeem KR (ed) *Crop Production and Global Environmental Issues*. pp 319-347.  
 1038 doi:10.1007/978-3-319-23162-4\_14  
 1039 Van Ooijen JW (2006) JoinMap 4, Software for the calculation of genetic linkage maps in experimental populations.  
 1040 Kyazma B.V., Wageningen, Netherlands.

1041 Van Ooijen JW (2009) MapQTL 6, Software for the mapping of quantitative trait loci in experimental populations  
1042 of diploid species. Kyazma B.V., Wageningen, Netherlands.

1043 Vogel KP, Tober D, Reece PE, Baltsensperger DD, Schuman G, Nicholson RA (2005) Registration of ‘Haymaker’  
1044 Intermediate Wheatgrass Registration by CSSA. Crop Sci 45:415-416. doi:10.2135/cropsci2005.0415

1045 Vu GT, Wicker T, Buchmann JP, Chandler PM, Matsumoto T, Graner A, Stein N (2010) Fine mapping and syntenic  
1046 integration of the semi-dwarfing gene *sdw3* of barley. Funct Integr Genomics 10:509-521.  
1047 doi:10.1007/s10142-010-0173-4

1048 Wagoner P (1990) Perennial grain new use for intermediate wheatgrass. Journal of Soil and Water Conservation  
1049 45:81-82

1050 Wang E et al. (2008) Control of rice grain-filling and yield by a gene with a potential signature of domestication.  
1051 Nat Genet 40:1370-1374. doi:10.1038/ng.220

1052 Wang GJ et al. (2014) Establishment and yield of perennial grass monocultures and binary mixtures for bioenergy in  
1053 North Dakota. Agron J 106:1605-1613. doi:10.2134/agronj14.0068

1054 Wang H et al. (2005) The origin of the naked grains of maize. Nature 436:714-719. doi:10.1038/nature03863

1055 Wang J, Liao X, Li Y, Zhou R, Yang X, Gao L, Jia J (2010) Fine mapping a domestication-related QTL for spike-  
1056 related traits in a synthetic wheat. Genome 53:798-804. doi:10.1139/g10-066

1057 Wang S et al. (2012) Control of grain size, shape and quality by *OsSPL16* in rice. Nat Genet 44:950-954.  
1058 doi:10.1038/ng.2327

1059 Wang Y et al. (2015) Copy number variation at the *GL7* locus contributes to grain size diversity in rice. Nat Genet  
1060 47:944-948. doi:10.1038/ng.3346

1061 Wendt T et al. (2016) *HvDep1* is a positive regulator of culm elongation and grain size in barley and impacts yield  
1062 in an environment-dependent manner. PLoS One 11. doi:10.1371/journal.pone.0168924

1063 Weng J et al. (2008) Isolation and initial characterization of *GW5*, a major QTL associated with rice grain width and  
1064 weight. Cell Res 18:1199-1209. doi:10.1038/cr.2008.307

1065 Win KT et al. (2017) A single base change explains the independent origin of and selection for the nonshattering  
1066 gene in African rice domestication. New Phytol 213:1925-1935. doi:10.1111/nph.14290

1067 Wu YZ, Fu YC, Zhao SS, Gu P, Zhu ZF, Sun CQ, Tan LB (2016) *CLUSTERED PRIMARY BRANCH 1*, a new allele  
1068 of *DWARF11*, controls panicle architecture and seed size in rice. Plant Biotechnol J 14:377-386.  
1069 doi:10.1111/pbi.12391

1070 Yan L, Loukoianov A, Tranquilli G, Helguera M, Fahima T, Dubcovsky J (2003) Positional cloning of the wheat  
1071 vernalization gene *VRN1*. Proc Natl Acad Sci USA 100:6263-6268. doi:10.1073/pnas.0937399100

1072 Yan LL et al. (2004) The wheat *VRN2* gene is a flowering repressor down-regulated by vernalization. Science  
1073 303:1640-1644. doi:10.1126/science.1094305

1074 Yan S et al. (2011) Seed size is determined by the combinations of the genes controlling different seed  
1075 characteristics in rice. Theor Appl Genet 123:1173-1181. doi:10.1007/s00122-011-1657-x

1076 Yang W-Y, Lu B-R, Hu X-R, Yu Y, Zhang Y (2005) Inheritance of the Triple-spikelet Character in a Tibetan  
1077 Landrace of Common Wheat. Genet Resour Crop Evol 52:847-851. doi:10.1007/s10722-003-6089-2

1078 Zadoks JC, Chang TT, Konzak CF (1974) A decimal code for the growth stages of cereals. *Weed Res* 14:415-421  
 1079 Zair W, Maxted N, Amri A (2018) Setting conservation priorities for crop wild relatives in the Fertile Crescent.  
 1080 *Genet Resour Crop Evol* 65:855-863. doi:10.1007/s10722-017-0576-3  
 1081 Zhang L, Zhao YL, Gao LF, Zhao GY, Zhou RH, Zhang BS, Jia JZ (2012a) *TaCKX6-D1*, the ortholog of rice  
 1082 *OsCKX2*, is associated with grain weight in hexaploid wheat. *New Phytol* 195:574-584.  
 1083 doi:10.1111/j.1469-8137.2012.04194.x  
 1084 Zhang X et al. (2017) Uncovering the genetic architecture of seed weight and size in intermediate wheatgrass  
 1085 through linkage and association mapping. *Plant Genome* 10:0. doi:10.3835/plantgenome2017.03.0022  
 1086 Zhang X et al. (2012b) Rare allele of *OsPPKL1* associated with grain length causes extra-large grain and a  
 1087 significant yield increase in rice. *Proc Natl Acad Sci USA* 109:21534-21539.  
 1088 doi:10.1073/pnas.1219776110  
 1089 Zhang XF et al. (2016) Establishment and optimization of genomic selection to accelerate the domestication and  
 1090 improvement of intermediate wheatgrass. *Plant Genome* 9. doi:10.3835/plantgenome2015.07.0059  
 1091 Zhang Y et al. (2015) Establishment of a 100-seed weight quantitative trait locus–allele matrix of the germplasm  
 1092 population for optimal recombination design in soybean breeding programmes. *J Exp Bot* 66:6311-6325.  
 1093 doi:10.1093/jxb/erv342  
 1094



**Figure caption**

**Figure 1.** Alignment of quantitative trait loci (QTLs), association-mapping markers, and possible candidate genes to 21 linkage groups (LG01 – LG21) and 21 chromosome sequences (CHR01 – CHR21) corresponding to seven homoeologous groups (HG1 – HG7) of allotetraploid ( $2n=6x=42$ ) intermediate wheatgrass. The 2-LOD drop-off intervals for 111 M26 x M35 QTLs (Tables 4 – 6) and 44 C3-2331 x C3-2595 (UMN) seed-size QTLs (Zhang et al. 2017) are indicated by filled and shaded box plots, respectively, on the right side of each linkage group. The physical position of UMN seed-size AM markers (Zhang et al. 2017) and possible candidate genes (Table 8) are shown in bold and italic text, respectively, on the left side of each chromosome. Graphed portions of each linkage group and chromosome sequence are scaled in centiMorgans (cM) in nucleotide megabases (MB), respectively, and normalized to comparable lengths.

**Table 1** Trait means with least significant differences (LSD) for cycle-3 (C3) grandparents and cycle-4 (C4) parents of the full-sib M26 x M36 population

Trait description	Trait	Units	C3 grandparents		C4 parents		LSD <sup>c</sup>	LSD levels
			C3_3471	C3_3941	M26 <sup>a</sup>	M35 <sup>b</sup>		
Seed mass	SEMA	mg	7.09 A	5.09 B	5.55 B	5.44 B	0.51	AB
Seed area	SEAR	mm <sup>2</sup>	5.56 A	4.50 C	4.56 C	4.79 B	0.18	ABC
Seed length	SELE	mm	5.29 A	4.22 D	4.61 B	4.36 C	0.11	ABCD
Seed width	SEWI	mm	1.35 C	1.38 B	1.27 D	1.43 A	0.03	ABCD
Seed shattering	SESH	%	5.0 C	30.0 B	28.8 B	59.4 A	0.6	ABC
Seed nakedness	SENA	%	94.1 A	28.7 C	71.3 B	30.3 C	2.7	ABC
Spiklets inflorescence <sup>-1</sup>	SPIN	no.	22.8 A	21.6 B	22.9 A	19.0 C	0.9	ABC
Florets spikelet <sup>-1</sup>	FLSP	no.	6.63 B	6.33 B	6.75 B	7.67 A	0.49	AB
Seeds floret <sup>-1</sup>	SEFL	no.	0.27 A	0.06 C	0.26 A	0.18 B	0.05	ABC
Seeds spikelet <sup>-1</sup>	SESP	no.	1.95 A	0.60 C	1.87 A	1.52 B	0.25	ABC
Seed yield inflorescence <sup>-1</sup>	SYIN	mg	323 A	72 D	237 B	149 C	21	ABCD
Inflorescence length	INLE	cm	28.0 B	27.6 B	30.3 A	26.0 C	1.2	ABC
Stem length	STLE	cm	114 B	128 A	128 A	124 A	6	AB
Stem width	STWI	mm	2.20 B	2.10 B	2.32 A	2.33 A	0.10	AB
Zadok's maturity	ZAMA	0-99	61.7 A	56.3 D	57.6 C	58.7 B	0.7	ABCD
Crown circumference	CRCI	cm	147 AB	145 AB	137 B	156 A	19	AB
Inflorescences crown <sup>-1</sup>	INCR	no.	219 A	197 AB	212 AB	184 B	31	AB

<sup>a</sup> Hybrid of C3\_3471 x C3\_3941 grandparents

<sup>b</sup> Hybrid of C3\_3941 x unknown grandparents

<sup>c</sup> Least significant differences,  $p \leq 0.05$

**Table 2** Trait means, relative standard deviations (RSD), ranges, and broad sense heritabilities with standard errors ( $H \pm SE$ ) for 266 M26 x M35 progeny across five location (KS or UT) x year (2013, 2014, and 2015) environments and overall average across five environments (AVG)

Trait	----- Means $\pm$ RSD (and ranges) -----						----- $H \pm SE$ -----	
	KS13	KS14	KS15	UT14	UT15	AVG	plot basis	genet basis
SEMA	4.57 $\pm$ 20% (2.43 - 8.95)	5.33 $\pm$ 14% (3.44 - 7.71)	5.16 $\pm$ 17% (2.09 - 8.70)	5.59 $\pm$ 17% (2.81 - 8.33)	5.27 $\pm$ 13% (3.27 - 7.83)	5.12 $\pm$ 13% (3.39 - 7.53)	0.43 $\pm$ 0.03	0.82 $\pm$ 0.02
SEAR	3.92 $\pm$ 12% (2.74 - 5.46)	5.58 $\pm$ 10% (4.22 - 7.21)	4.70 $\pm$ 11% (3.05 - 6.28)	5.32 $\pm$ 11% (3.66 - 7.42)	4.47 $\pm$ 10% (3.33 - 6.26)	4.60 $\pm$ 9% (3.41 - 6.09)	0.41 $\pm$ 0.03	0.80 $\pm$ 0.02
SELE	4.22 $\pm$ 7% (3.56 - 4.99)	5.04 $\pm$ 6% (4.20 - 5.92)	5.01 $\pm$ 7% (4.22 - 5.88)	4.79 $\pm$ 6% (4.07 - 5.71)	4.18 $\pm$ 6% (3.51 - 4.90)	4.47 $\pm$ 6% (3.87 - 5.21)	0.54 $\pm$ 0.03	0.87 $\pm$ 0.02
SEWI	1.22 $\pm$ 7% (1.02 - 1.46)	1.45 $\pm$ 6% (1.20 - 1.67)	1.38 $\pm$ 7% (1.01 - 1.59)	1.45 $\pm$ 6% (1.18 - 1.68)	1.35 $\pm$ 5% (1.17 - 1.58)	1.34 $\pm$ 5% (1.12 - 1.51)	0.42 $\pm$ 0.03	0.82 $\pm$ 0.02
SESH	---	29.7 $\pm$ 70% (0.0 - 84.7)	40.0 $\pm$ 49% (0.0 - 87.2)	51.8 $\pm$ 53% (0.0 - 86.8)	12.9 $\pm$ 78% (0.0 - 52.0)	33.7 $\pm$ 52% (0.0 - 67.5)	0.38 $\pm$ 0.03	0.71 $\pm$ 0.3
SENA	67.5 $\pm$ 20% (0.0 - 100.0)	29.6 $\pm$ 83% (0.0 - 87.0)	10.1 $\pm$ 94% (0.0 - 55.0)	77.5 $\pm$ 19% (21.7 - 98.0)	66.1 $\pm$ 17% (26.9 - 92.3)	55.3 $\pm$ 20% (23.1 - 83.5)	0.30 $\pm$ 0.03	0.71 $\pm$ 0.03
SPIN	19.9 $\pm$ 11% (8.1 - 26.6)	20.8 $\pm$ 13% (10.6 - 28.3)	21.8 $\pm$ 11% (14.3 - 28.6)	19.9 $\pm$ 15% (11.6 - 37.1)	18.0 $\pm$ 13% (8.5 - 24.5)	19.7 $\pm$ 11% (11.2 - 29.2)	0.30 $\pm$ 0.03	0.75 $\pm$ 0.02
FLSP	8.12 $\pm$ 17% (3.51 - 12.5)	6.72 $\pm$ 13% (4.51 - 10.2)	6.86 $\pm$ 11% (5.01 - 8.51)	---	---	7.24 $\pm$ 11% (4.68 - 9.92)	0.22 $\pm$ 0.03	0.56 $\pm$ 0.03
SEFL	0.12 $\pm$ 47% (0.00 - 0.47)	0.10 $\pm$ 71% (0.00 - 0.31)	0.22 $\pm$ 44% (0.00 - 0.49)	---	---	0.15 $\pm$ 47% (0.00 - 0.33)	0.43 $\pm$ 0.03	0.77 $\pm$ 0.03
SESP	0.98 $\pm$ 79% (0.00 - 4.4)	0.63 $\pm$ 56% (0.00 - 1.8)	1.46 $\pm$ 44% (0.00 - 3.49)	2.08 $\pm$ 33% (0.03 - 3.70)	1.16 $\pm$ 43% (0.00 - 3.10)	1.17 $\pm$ 43% (0.00 - 2.32)	0.44 $\pm$ 0.03	0.77 $\pm$ 0.02
SYIN	95 $\pm$ 86% (0 - 454)	75 $\pm$ 64% (2 - 247)	170 $\pm$ 48% (2 - 437)	235 $\pm$ 38% (2 - 525)	111 $\pm$ 49% (0 - 274)	120 $\pm$ 53% (0 - 326)	0.50 $\pm$ 0.03	0.84 $\pm$ 0.02
INLE	---	28.4 $\pm$ 15% (18.8 - 41.8)	31.0 $\pm$ 12% (22.8 - 41.8)	25.6 $\pm$ 12% (18.2 - 35.4)	23.4 $\pm$ 14% (16.2 - 33.0)	27.3 $\pm$ 11% (20.1 - 37.3)	0.56 $\pm$ 0.03	0.86 $\pm$ 0.01
STLE	41 $\pm$ 10% (30 - 53)	132 $\pm$ 7% (93 - 153)	141 $\pm$ 6% (108 - 165)	170 $\pm$ 11% (76 - 200)	182 $\pm$ 9% (117 - 219)	121 $\pm$ 11% (26 - 149)	0.47 $\pm$ 0.03	0.82 $\pm$ 0.02
STWI	2.43 $\pm$ 14% (1.33 - 3.34)	1.97 $\pm$ 15% (1.27 - 4.50)	2.18 $\pm$ 13% (1.37 - 2.84)	2.09 $\pm$ 10% (1.51 - 2.68)	1.67 $\pm$ 13% (0.91 - 2.30)	2.10 $\pm$ 10% (1.39 - 2.71)	0.33 $\pm$ 0.03	0.75 $\pm$ 0.02
ZAMA	65.8 $\pm$ 7% (32.0 - 71.9)	59.3 $\pm$ 5% (52.4 - 68.9)	58.7 $\pm$ 4% (53.1 - 65.9)	52.3 $\pm$ 4% (48.1 - 57.0)	47.1 $\pm$ 8% (38.8 - 57.9)	56.6 $\pm$ 5% (44.7 - 63.2)	0.48 $\pm$ 0.03	0.82 $\pm$ 0.02
CRCI	---	131 $\pm$ 15% (81 - 179)	---	127 $\pm$ 23% (59 - 234)	165 $\pm$ 18% (53 - 256)	138 $\pm$ 16% (80 - 196)	0.24 $\pm$ 0.03	0.58 $\pm$ 0.04
INCR	---	160 $\pm$ 24% (37 - 299)	180 $\pm$ 25% (60 - 324)	121 $\pm$ 35% (8 - 280)	244 $\pm$ 35% (7 - 563)	176 $\pm$ 24% (43 - 322)	0.23 $\pm$ 0.03	0.63 $\pm$ 0.03

**Table 3** Trait correlations among parents and progeny of M26 x M35 family

	SEMA	SEAR	SELE	SEWI	SESH	SENA	SPIN	FLSP	SEFL	SESP	SYIN	INLE	STLE	STWI	ZAMA	CRCI	INCR
SEMA	.	***	***	***	***	---	---	***	***	***	***	***	***	***	***	---	---
SEAR	0.67	.	***	***	***	---	---	***	***	***	***	***	***	***	***	***	---
SELE	0.65	0.86	.	***	***	---	---	***	***	***	***	***	*	***	***	***	---
SEWI	0.52	0.8	0.42	.	***	***	---	***	***	***	***	***	***	***	***	---	---
SESH	0.24	0.24	0.17	0.26	.	***	---	**	***	***	***	***	***	***	***	*	***
SENA	---	---	---	-0.13	-0.2	.	---	**	---	---	---	---	---	---	---	---	---
SPIN	---	---	---	---	---	---	.	***	---	---	***	***	***	***	***	---	*
FLSP	0.16	0.22	0.13	0.26	0.13	-0.12	0.24	.	---	***	***	***	***	***	***	***	***
SEFL	0.2	0.22	0.23	0.14	0.38	---	---	---	.	***	***	---	***	***	***	---	***
SESP	0.15	0.2	0.16	0.17	0.37	---	---	0.23	0.95	.	***	*	***	***	***	---	***
SYIN	0.44	0.4	0.38	0.29	0.38	---	0.27	0.26	0.82	0.87	.	***	***	***	***	---	***
INLE	0.23	0.27	0.28	0.18	0.11	---	0.44	0.38	---	0.08	0.27	.	***	***	---	---	***
STLE	0.22	0.18	0.08	0.22	0.36	---	0.17	0.17	0.12	0.21	0.26	0.23	.	***	***	***	---
STWI	0.27	0.3	0.21	0.31	0.21	---	0.33	0.49	0.18	0.29	0.39	0.48	0.21	.	***	---	---
ZAMA	0.21	0.15	0.12	0.13	0.28	---	0.11	0.27	0.34	0.32	0.35	---	0.25	0.27	.	***	***
CRCI	---	-0.11	-0.11	---	0.09	---	---	-0.22	---	---	---	---	0.23	---	0.16	.	***
INCR	---	---	---	---	0.14	---	0.08	-0.2	0.21	0.12	0.11	-0.1	0.3	---	0.23	0.55	.

\*, \*\*, and \*\*\* Significant at  $P \leq 0.05$ ,  $P \leq 0.01$  and  $P \leq 0.001$  controlling for multiple tests with Bonferroni correction, respectively

<sup>a</sup> Based on standard normal deviations of SENA and SESH.

**Table 4** Linkage groups (LGs) with significant QTL effects in M26 x M35 population across five location (KS or UT) x year (2013, 2014, and 2015) environments and overall average across five environments (AVG)

Trait	KS13 (LG)	KS14 (LG)	KS15 (LG)	UT14 (LG)	UT15 (LG)	AVG (LG)	Total number of QTLs
SEMA	1*	15*	---	6*	---	6, 11*	5 (4*)
SEAR	6*, 20	7	7*	9	17*,	7, 9*, 17, 20*, 19*	11 (6*)
SELE	6	6, 14*, 21	6, 21, 19	6, 9*	6, 17*, 20*, 21*	6*, 17, 20, 21, 19*	18 (7*)
SEWI	6*, 20	9*, 15*	11*,	8*, 16	15 17*	11, 16*, 20*	12 (8*)
SESH	---	11, 12*	6*, 11	11	14*	10*, 11*	8 (5*)
SENA	2, 15*	2, 10*, 11*, 14*, 15	2*, 10, 11	6, 10	2, 6, 12*	2, 6*, 5*, 10, 18*	20 (9*)
SPIN	3, 6*, 7	10*	3, 10, 13	---	4*, 7, 9*, 14*, 13*	3*, 7*	14 (8*)
FLSP	---	---	9*	---	---	4*, 8*, 9, 16*	5 (4*)
SEFL	6, 12*, 14*, 13*, 15 *	6	9*, 10*, 11*	---	---	6*, 9	11 (8*)
SESP	6, 14*, 15*	6*	1*, 6, 7, 10*, 11*	7*, 10	6, 7, 14, 13*	7, 10, 14, 15	19 (8*)
SYIN	6, 12*, 15*	6*, 13*, 21*	1*	4*	---	3*, 6, 4	11 (8*)
INLE	---	9, 12, 13	9, 14*	2*, 9, 12*, 14	5*, 7*, 9, 12, 14, 13	9*, 12, 14, 13*	19 (7*)
STLE	19*,	5*, 9 16	16*	5, 10 13*	9*, 10*, 11*, 16	5, 8*	14 (8*)
STWI	15*	---	6, 9	6, 9	4	6*, 4*, 9*	9 (4*)
ZAMA	15	6, 10*, 14*, 18*, 11, 3	14, 11, 15*	6, 11	6*, 11, 17*, 16*, 3*	6, 10, 14, 11*, 3, 12*, 15	24 (10*)
CRCI	---	---	---	12*, 13	---	13*, 19*	4 (3*)
INCR	---	---	---	10*, 13*, 17*	15*	10, 15	6 (4*)
Total number of QTLs	25 (15*)	32 (17*)	31 (15*)	28 (12*)	37 (22*)	57 (30*)	210 (111*)

\* Most significant QTL for each linkage group by trait combination.

**Table 5** Peak LOD scores for each linkage group (LG) and homoeologous group (HG) of M26 x M36 family

LG	HG	SEMA	SEAR	SELE	SEWI	SESH	SENA	SPIN	FLSP	SEFL	SESP	SYIN	INLE	STLE	STWI	ZAMA	CRCI	INCR
1	1	3.5 <sup>b</sup>	---	---	---	---	---	---	---	---	5.3 <sup>c</sup>	5.7 <sup>c</sup>	---	---	---	---	---	---
2	1	---	---	---	---	---	8.0 <sup>c</sup>	---	---	---	---	---	5.7 <sup>c</sup>	---	---	---	---	---
3	1	---	---	---	---	---	---	5.2 <sup>c</sup>	---	---	---	3.5 <sup>a</sup>	---	---	---	6.8 <sup>b</sup>	---	---
4	2	---	---	---	---	---	---	5.8 <sup>c</sup>	6.1 <sup>c</sup>	---	---	5.6 <sup>c</sup>	---	---	6.5 <sup>c</sup>	---	---	---
5	2	---	---	---	---	---	6.6 <sup>c</sup>	---	---	---	---	---	5.6	6.3	---	---	---	---
6	2	6.8 <sup>c</sup>	6.0 <sup>c</sup>	12.0 <sup>c</sup>	5.1 <sup>c</sup>	5.0 <sup>c</sup>	5.6 <sup>a</sup>	5.6 <sup>c</sup>	---	8.1 <sup>c</sup>	6.6 <sup>c</sup>	8.4 <sup>c</sup>	---	---	5.1 <sup>c</sup>	5.8 <sup>c</sup>	---	---
7	3	---	5.2 <sup>c</sup>	---	---	---	---	6.6 <sup>a</sup>	---	---	5.3 <sup>b</sup>	---	3.6 <sup>a</sup>	---	---	---	---	---
8	3	---	---	---	7.2 <sup>c</sup>	---	---	---	6.2 <sup>c</sup>	---	---	---	---	5.9 <sup>c</sup>	---	---	---	---
9	3	---	5.0 <sup>c</sup>	4.8 <sup>c</sup>	3.7 <sup>a</sup>	---	---	4.2 <sup>b</sup>	7.5 <sup>c</sup>	6.5 <sup>c</sup>	---	---	13.0 <sup>c</sup>	5.7 <sup>b</sup>	4.4 <sup>b</sup>	---	---	---
10	4	---	---	---	---	3.9 <sup>a</sup>	7.6 <sup>c</sup>	6.4 <sup>c</sup>	---	5.0 <sup>c</sup>	5.8 <sup>c</sup>	---	---	9.4 <sup>a</sup>	---	6.3 <sup>c</sup>	---	7.2 <sup>c</sup>
11	4	3.8 <sup>b</sup>	---	---	6.7 <sup>c</sup>	15.0 <sup>c</sup>	6.9 <sup>c</sup>	---	---	3.9 <sup>a</sup>	5.3 <sup>c</sup>	---	---	6.0 <sup>c</sup>	---	7.8 <sup>c</sup>	---	---
12	4	---	---	---	---	4.0 <sup>b</sup>	5.5 <sup>c</sup>	---	---	5.1 <sup>c</sup>	---	3.8 <sup>b</sup>	5.3 <sup>c</sup>	---	---	5.5 <sup>c</sup>	3.8 <sup>b</sup>	---
13	5	---	---	---	---	---	---	5.3 <sup>a</sup>	---	3.6 <sup>b</sup>	3.7 <sup>b</sup>	3.9 <sup>a</sup>	5.6 <sup>c</sup>	5.6 <sup>c</sup>	---	---	6.3 <sup>c</sup>	5.3 <sup>c</sup>
14	5	---	---	5.1 <sup>b</sup>	---	3.7 <sup>b</sup>	5.0 <sup>c</sup>	5.5 <sup>c</sup>	---	6.9 <sup>c</sup>	6.6 <sup>c</sup>	---	5.4 <sup>c</sup>	---	---	5.9 <sup>c</sup>	---	---
15	5	4.5 <sup>a</sup>	---	---	5.4 <sup>a</sup>	---	5.4 <sup>c</sup>	---	---	6.0 <sup>c</sup>	5.7 <sup>c</sup>	7.1 <sup>c</sup>	---	---	3.6 <sup>b</sup>	7.4 <sup>c</sup>	---	4.1 <sup>b</sup>
16	6	---	---	---	4.6 <sup>a</sup>	---	---	---	---	---	---	---	---	5.2 <sup>a</sup>	---	6.3 <sup>c</sup>	---	---
17	6	---	6.0 <sup>c</sup>	4.2 <sup>b</sup>	3.9 <sup>b</sup>	---	---	---	---	---	---	---	---	---	---	6.0 <sup>c</sup>	---	3.7 <sup>b</sup>
18	6	---	---	---	---	---	3.5 <sup>a</sup>	---	---	---	---	---	---	---	---	6.8 <sup>c</sup>	---	---
19	7	---	4.2 <sup>a</sup>	4.0 <sup>a</sup>	---	---	---	---	5.6 <sup>c</sup>	---	---	---	---	4.4 <sup>a</sup>	---	---	6.0 <sup>c</sup>	---
20	7	---	5.1 <sup>b</sup>	5.4 <sup>c</sup>	4.9 <sup>c</sup>	---	---	---	---	---	---	---	---	---	---	---	---	---
21	7	---	---	6.3 <sup>c</sup>	---	---	---	---	---	---	---	3.6 <sup>a</sup>	---	---	---	---	---	---
M26 <sup>a</sup>	1	1	1	3	1	2	2	0	1	0	3	1	3	0	0	0	0	0
M35 <sup>b</sup>	2	1	2	1	2	0	1	0	1	2	1	0	1	2	1	1	1	2
CP <sup>c</sup>	1	4	4	4	2	7	5	4	6	6	4	6	4	2	9	2	2	2
Total	4	6	7	8	5	9	8	4	8	8	8	8	7	8	4	10	3	4

<sup>a</sup> M26 two-way pseudo-tester (TWPT) QTL<sup>b</sup> M35 TWPT QTL<sup>c</sup> Cross-pollinators (CP) QTL

**Table 6** Estimates of maximum QTL effects ( $\alpha, \gamma, \tau$ ) based on six QTL analyses of the average trait values within or among five environments (Tables 4 and 5), total magnitudes of maximum (Max) QTL effects, and total magnitudes of average (Avg) QTL effects over five environments reported as a percentage of population mean where  $\alpha$  is difference between QTL alleles of the M26 parent,  $\gamma$  is difference between QTL alleles of the M35 parent, and  $\tau$  is intralocus interaction

LG	HG	SEMA	SEAR	SELE	SEWI	SESH	SENA	SPIN	FLSP	SEFL	SESP	SYIN	INLE	STLE	STWI	ZAMA	CRCI	INCR
1	1	0 <sup>a</sup> ,5,0 <sup>a</sup>	---	---	---	---	---	---	---	---	13,8,3	10,6,1	---	---	---	---	---	---
2	1	---	---	---	---	---	-21,-11,-7	---	---	---	---	---	-3,-1,0	---	---	---	---	---
3	1	---	---	---	---	---	---	3,0,1	---	---	---	-12,0 <sup>a</sup> ,0 <sup>a</sup>	---	---	---	0 <sup>a</sup> ,3,0 <sup>a</sup>	---	---
4	2	---	---	---	---	---	---	-1,2,-3	-2,2,-2	---	---	-5,-1,-11	---	---	-2,2,-2	---	---	---
5	2	---	---	---	---	---	-1,3,3	---	---	---	---	---	-3,-3,-1	-1,-2,1	---	---	---	---
6	2	-2,-2,-3	-4,0,-3	-2,-1,0	-1,1,-2	-7,-7,-14	-5,0 <sup>a</sup> ,0 <sup>a</sup>	-2,3,-3	---	-9,-9,-13	-10,-10,-15	-12,-12,-15	---	---	-2,-2,-1	-2,-1,-2	---	---
7	3	---	-3,0,0	---	---	---	---	3,0 <sup>a</sup> ,0 <sup>a</sup>	---	---	0 <sup>a</sup> , -10,0 <sup>a</sup>	---	3,0 <sup>a</sup> ,0 <sup>a</sup>	---	---	---	---	---
8	3	---	---	---	0,0,2	---	---	---	-1,5,-3	---	---	---	---	-2,0,2	---	---	---	---
9	3	---	-2,1,-1	-2,0,0	-1,0 <sup>a</sup> ,0 <sup>a</sup>	---	---	0 <sup>a</sup> ,3,0 <sup>a</sup>	-2,2,-2	9,-8,6	---	---	-3,5,0	0 <sup>a</sup> ,2,0 <sup>a</sup>	0 <sup>a</sup> ,3,0 <sup>a</sup>	---	---	---
10	4	---	---	---	---	6,0 <sup>a</sup> ,0 <sup>a</sup>	-10,10,3	0,4,-2	---	2,12,-5	5,14,-7	---	---	-3,0 <sup>a</sup> ,0 <sup>a</sup>	---	-1,1,-1	---	0,9,-1
11	4	0 <sup>a</sup> ,3,0 <sup>a</sup>	---	---	2,2,0	17,7,5	-16,-4,-2	---	---	-12,0 <sup>a</sup> ,0 <sup>a</sup>	13,7,2	---	---	2,0,2	---	1,0,-1	---	---
12	4	---	---	---	---	0 <sup>a</sup> , -16,0 <sup>a</sup>	2,4,3	---	---	-8,-8,-3	---	0 <sup>a</sup> , -22,0 <sup>a</sup>	1,-3,0	---	---	-1,0,-1	0 <sup>a</sup> ,6,0 <sup>a</sup>	---
13	5	---	---	---	---	---	---	3,0 <sup>a</sup> ,0 <sup>a</sup>	---	0 <sup>a</sup> ,19,0 <sup>a</sup>	0 <sup>a</sup> ,10,0 <sup>a</sup>	17,0 <sup>a</sup> ,0 <sup>a</sup>	3,1,-1	-2,0,-2	---	---	-2,-1,-4	-6,1,-5
14	5	---	---	0 <sup>a</sup> ,2,0 <sup>a</sup>	---	0 <sup>a</sup> ,3,0 <sup>a</sup>	-3,-14,3	1,3,0	---	-5,-15,-1	-6,-16,-4	---	1,4,0	---	---	0,-1,0	---	---
15	5	4,0 <sup>a</sup> ,0 <sup>a</sup>	---	---	2,0 <sup>a</sup> ,0 <sup>a</sup>	---	-5,1,1	---	---	-2,-14,7	-2,-15,8	-1,-14,9	---	---	0 <sup>a</sup> , -4,0 <sup>a</sup>	0,-1,0	---	0 <sup>a</sup> , -9, $\tau$ <sup>a</sup>
16	6	---	---	---	-1,0 <sup>a</sup> ,0 <sup>a</sup>	---	---	---	-3,1,1	---	---	---	---	2,0 <sup>a</sup> ,0 <sup>a</sup>	---	-2,2,0	---	---
17	6	---	2,-3,1	0 <sup>a</sup> , -2,0 <sup>a</sup>	0 <sup>a</sup> , -1,0 <sup>a</sup>	---	---	---	---	---	---	---	---	---	---	-1,2,1	---	0 <sup>a</sup> , -8, $\tau$ <sup>a</sup>
18	6	---	---	---	---	---	-4,0 <sup>a</sup> ,0 <sup>a</sup>	---	---	---	---	---	---	---	---	0,-1,0	---	---
19	7	---	2,0 <sup>a</sup> ,0 <sup>a</sup>	1,0 <sup>a</sup> ,0 <sup>a</sup>	---	---	---	---	---	---	---	---	---	3,0 <sup>a</sup> ,0 <sup>a</sup>	---	---	3,2,-3	---
20	7	---	0 <sup>a</sup> ,3,0 <sup>a</sup>	0,2,0	0,1,1	---	---	---	---	---	---	---	---	---	---	---	---	---
21	7	---	---	-1,2,-1	---	---	---	---	---	---	---	-17,0 <sup>a</sup> ,0 <sup>a</sup>	---	---	---	---	---	---
Max		7,11,3	13,6,6	7,8,1	8,5,5	29,33,19	67,47,22	14,16,9	8,10,7	46,86,37	48,90,39	73,55,37	17,17,3	15,5,8	3,10,3	7,12,6	5,10,8	6,27,6
(total)		(21)	(25)	(16)	(18)	(81)	(136)	(39)	(25)	(169)	(177)	(164)	(36)	(27)	(17)	(25)	(22)	(40)
Avg		5,8,3	8,5,2	6,4,1	5,5,3	30,26,12	28,12,11	14,12,6	7,11,6	33,50,31	32,48,38	32,27,24	14,14,4	8,5,6	3,7,3	5,6,3	5,8,8	2,15,7
(total)		(15)	(16)	(12)	(12)	(68)	(52)	(31)	(24)	(114)	(118)	(82)	(32)	(19)	(14)	(15)	(20)	(24)

<sup>a</sup> Non-estimable  $\alpha$ ,  $\gamma$ , or  $\tau$  effect for QTLs detected using two-way pseudo-testcross model

**Table 7** Total percent variation explained (PVE) by M26 x M35 QTL markers using a two-way pseudo-testcross (TWPT) or a cross-pollinators (CP) across five location (KS or UT) x year (2013, 2014, and 2015) environments and overall averages across five environments (AVG) including AVGs by parent (M26 or M35)

Trait	Model	Number markers	KS13 PVE	KS14 PVE	KS15 PVE	UT14 PVE	UT15 PVE	AVG PVE	AVG M26 PVE	AVG M35 PVE
SEMA	TWPT	5	14.3	16.2	8.0	11.0	11.3	15.1	4.8	11.5
	CP	4	16.6	19.6	12.7	18.1	12.9	20.8	---	---
SEAR	TWPT	10	18.1	21.4	22.7	24.2	17.9	31.5	21.1	15.6
	CP	6	28.1	22.8	18.8	23.9	18.4	29.3	---	---
SELE	TWPT	11	28.4	35.7	36.4	33.3	28.0	39.8	26.6	21.0
	CP	7	29.8	34.7	29.5	31.7	27.7	37.5	---	---
SEWI	TWPT	13	17.6	26.4	19.4	21.1	27.1	28.6	15.2	18.2
	CP	8	27.9	30.0	25.2	27.3	29.6	33.9	---	---
SESH	TWPT	7	---	28.74	16.4	26.4	16.1	29.6	16.8	15.8
	CP	5	---	35.4	26.5	25.7	16.2	29.5	---	---
SENA	TWPT	16	26.3	47.3	46.5	26.3	26.1	44.8	35.4	17.9
	CP	9	28.8	47.7	40.9	28.8	34.7	46.8	---	---
SPIN	TWPT	13	14.4	25.0	22.6	12.3	36.5	27.1	22.6	9.1
	CP	8	22.5	35.6	24.9	13.5	38.2	31.7	---	---
FLSP	TWPT	8	17.8	18.9	26.0	---	---	28.3	16.8	12.0
	CP	4	16.6	16.8	25.3	---	---	27.4	---	---
SEFL	TWPT	14	33.1	15.5	36.4	---	---	34.1	15.2	23.9
	CP	8	35.5	20.2	33.9	---	---	36.8	---	---
SESP	TWPT	15	28.2	17.4	30.1	25.7	32.5	29.7	10.6	20.8
	CP	8	32.7	22.7	29.0	23.6	31.0	30.1	---	---
SYIN	TWPT	12	27.7	31.3	24.3	20.5	22.7	26.9	14.3	13.9
	CP	8	32.7	39.1	26.9	21.9	21.3	27.8	---	---
INLE	TWPT	13	---	31.2	27.0	38.0	32.6	36.9	16.1	25.5
	CP	7	---	30.3	24.7	33.6	34.4	35.4	---	---
STLE	TWPT	12	17.7	25.4	28.6	23.5	37.2	23.2	14.5	12.5
	CP	8	26.3	27.3	34.9	30.5	28.9	38.0	---	---
STWI	TWPT	6	19.7	13.3	19.5	21.5	16.2	25.5	12.5	15.8
	CP	4	22.6	12.3	20.4	20.9	18.7	25.4	---	---
ZAMA	TWPT	19	27.6	41.4	40.4	42.74	44.5	45.2	19.5	30.4
	CP	10	34.2	49.2	45.3	44.0	46.7	49.6	---	---
CRCI	TWPT	5	---	9.6	---	8.7	8.9	8.4	1.9	6.8
	CP	3	---	14.7	---	14.2	13.6	19.2	---	---
INCR	TWPT	6	---	6.5	6.6	22.6	9.9	13.8	3.3	11.6
	CP	4	---	8.5	6.2	24.6	7.6	14.5	---	---







**Table 8** Description of 42 candidate genes located in 87 different chromosome regions containing one or more relevant quantitative trait loci (QTL) identified in the intermediate wheatgrass M26 x M35 or C3\_2332 x C3\_2595 (UMN) families, listed by homoeologous group (HG)

HG	Candidate gene	Chromosome (QTL)
1	<i>Barren inflorescence 4 (BIF4)</i>	CHR03 (SPIN)
1	<i>Grain weight Chr5 QTL (GW5)</i>	CHR01 (SEMA), CHR02 (SEMA, SEAR, SELE, SEWI), CHR03 (SELE)
1	<i>Seedstick (STK)</i>	CHR02 <sup>a</sup> (SENA)
1	<i>Self-incompatibility (S)</i>	CHR01 (SEMA), CHR02 (SEMA, SEAR, SELE, SEWI)
2	<i>Clustered primary branch 1 (CPB1)</i>	CHR05 (STLE), CHR06 (SEMA, SEAR, SELE, SEWI, SYIN)
2	<i>Grain incomplete filling (GIF1)</i>	CHR04 (SEAR, SELE, SEWI, SYIN), CHR06 (SEMA, SEAR, SELE, SEWI, SYIN)
2	<i>Grain length Chr7 QTL (GL7)</i>	CHR04 (SEAR, SELE, SEWI, SYIN), CHR06 (SEMA, SEAR, SELE, SEWI, SYIN)
2	<i>Self-incompatibility (Z)</i>	CHR01 <sup>b</sup> (SESP, SYIN), CHR06 (SEMA, SEAR, SELE, SEWI, SEFL, SESP, SYIN)
2	<i>Semidwarf 3 (SDW3)</i>	CHR05 (STLE)
2	<i>Six-rowed spike 1 (VRS1)</i>	CHR04 (SPIN, FLSP), CHR06 <sup>a</sup> (SEFL, SESP)
2	<i>Sucrose synthase 2 (SUS2)</i>	CHR04 (SEAR, SELE, SEWI, SYIN), CHR06 (SEMA, SEAR, SELE, SEWI, SYIN)
2	<i>Xylogalacturonan deficient (XGD)</i>	CHR04 (SEAR, SELE, SEWI, SYIN), CHR06 (SEMA, SEAR, SELE, SEWI, SYIN)
3	<i>Barren inflorescence 1 (BIF1)</i>	CHR07 (SPIN), CHR09 (SPIN, FLSP)
3	<i>Barren stalk 1 (BA1)</i>	CHR07 (SPIN), CHR09 (SPIN, FLSP)
3	<i>Brassinosteroid-insensitive 1 (BRI1)</i>	CHR07 (INLE), CHR09 (INLE, STLE)
3	<i>Grain number Chr1 QTL (GN1)</i>	CHR07 (SEAR, SPIN), CHR08 <sup>a</sup> (SEWI)
3	<i>Grain size Chr5 QTL (GS5)</i>	CHR07 (SEAR), CHR08 (SEWI), CHR09 (SEAR, SEWI)
3	<i>Semidwarf 1 (DENSO)</i>	CHR08 (STLE), CHR09 (STLE)
3	<i>Six-rowed spike 4 (VRS4)</i>	CHR07 (SPIN), CHR09 (SPIN)
4	<i>Phytochrome A (PHYA)</i>	CHR10 (ZAMA)
4	<i>Phytochrome B (PHYB)</i>	CHR10 (ZAMA), CHR11 (ZAMA)
4	<i>Reduced height (RHT)</i>	CHR10 (STLE)
4	<i>Seuss (SEU)</i>	CHR10 <sup>a</sup> (SESH, SENA), CHR12 <sup>a</sup> (SESH)
5	<i>Barren inflorescence 2 (BIF2)</i>	CHR14 (SPIN)
5	<i>Constans 3 (CO3)</i>	CHR14 (ZAMA), CHR15 (ZAMA)
5	<i>Dense and erect panicle 1 (DEP1)</i>	CHR13 (INLE, SPIN), CHR14 (INLE, SPIN)
5	<i>Grain length Chr3 QTL (GL3)</i>	CHR13 <sup>a</sup> (SEWI), CHR14 <sup>a</sup> (SEMA, SEAR, SELE), CHR15 <sup>a</sup> (SEMA, SEAR, SELE, SEWI)
5	<i>Phytochrome C (PHYC)</i>	CHR14 (ZAMA)
5	<i>Q free-threshing (Q)</i>	CHR14 (INLE, SESH, SENA), CHR15 (SENA)
5	<i>Vernalization 1 (VRN1)</i>	CHR14 (ZAMA)
6	<i>Constans 2 (CO2)</i>	CHR16 (ZAMA), CHR17 (ZAMA)
6	<i>Constans 5 (CO5)</i>	CHR17 (ZAMA)
6	<i>Constans 7 (CO7)</i>	CHR16 (ZAMA), CHR17 (ZAMA)
6	<i>GDSL lipase (GLIP)</i>	CHR16 (SEWI), CHR17 (SEMA, SEAR, SELE, SEWI), CHR18 (SEAR)
6	<i>Grain weight Chr2 QTL (GW2)</i>	CHR16 (SEWI), CHR17 (SEMA, SEAR, SELE, SEWI), CHR18 (SEAR)
6	<i>Growth-regulating factor 4 (GRF4)</i>	CHR16 (SEWI), CHR17 (SEMA, SEAR, SELE, SEWI), CHR18 (SEAR, SENA)
7	<i>Fertilization independent seed (FEI)</i>	CHR19 <sup>a</sup> (SEAR, SELE), CHR20 (SEMA, SEAR, SELE, SEWI), CHR21 (SELE)
7	<i>Glucan, water-dikinase (GWD)</i>	CHR19 (SEAR, SELE), CHR20 (SEAR, SELE), CHR21 (SELE)
7	<i>Grain width Chr8 QTL (GW8)</i>	CHR19 (SEAR, SELE), CHR20 (SEAR, SELE), CHR21 (SELE)
7	<i>Sucrose synthase 1 (SUS1)</i>	CHR19 <sup>a</sup> (SEAR, SELE), CHR20 <sup>a</sup> (SEMA, SEAR, SELE, SEWI), CHR21 (SELE)
7	<i>Thousand-grain weight 6 (TGW6)</i>	CHR15 <sup>a, b</sup> (SEMA, SEAR, SELE, SEWI), CHR19 (SEAR, SELE), CHR20 (SEAR, SELE, SEWI), CHR21 (SELE, SYIN)
7	<i>Thousand-grain weight Chr7A (TGW7A)</i>	CHR19 (SEAR, SELE), CHR20 (SEAR, SELE), CHR21 (SELE)

<sup>a</sup> Multiple gene loci present on same chromosome

<sup>b</sup> Paralogous gene locus






HG7


CHR19

LG19



CHR20

LG20



CHR21

LG21

